



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/44, C07D 401/14, 405/14, 409/14</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/56377</b> <b>(43) International Publication Date:</b> 17 December 1998 (17.12.98)
<b>(21) International Application Number:</b> PCT/US98/12387 <b>(22) International Filing Date:</b> 12 June 1998 (12.06.98) <b>(30) Priority Data:</b> 60/050,904                      13 June 1997 (13.06.97)                      US <b>(71) Applicant (for all designated States except US):</b> SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ADAMS, Jerry, L. [US/US]; 611 Forest Road, Wayne, PA 19087 (US). GALLAGHER, Timothy [US/US]; 255 Manor Road, Harleysville, PA 19438 (US). OSIFO, Irennegbe, Kelly [US/US]; 309 Conestoga Way, Eaglesville, PA 19403 (US). <b>(74) Agents:</b> DINNER, Dara, L. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).		<b>(81) Designated States:</b> AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> NOVEL PYRAZOLE AND PYRAZOLINE SUBSTITUTED COMPOUNDS		
<b>(57) Abstract</b>		
<p>Novel pyridyl or pyrimidinyl substituted pyrazole and pyrazoline compounds and compositions for use in therapy.</p>		

***FOR THE PURPOSES OF INFORMATION ONLY***

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## NOVEL PYRAZOLE AND PYRAZOLINE SUBSTITUTED COMPOUNDS

### FIELD OF THE INVENTION

This invention relates to a novel group of pyrazole containing compounds, processes for the preparation thereof, the use thereof in treating cytokine mediated diseases and pharmaceutical compositions for use in such therapy.

### BACKGROUND OF THE INVENTION

Intracellular signal transduction is the means by which cells respond to extracellular stimuli. Regardless of the nature of the cell surface receptor (e. g. protein tyrosine kinase or seven-transmembrane G-protein coupled), protein kinases and phosphatases along with phospholipases are the essential machinery by which the signal is further transmitted within the cell [Marshall, J. C. Cell, **80**, 179-278 (1995)]. Protein kinases can be categorized into five classes with the two major classes being, tyrosine kinases and serine / threonine kinases depending upon whether the enzyme phosphorylates its substrate(s) on specific tyrosine(s) or serine / threonine(s) residues [Hunter, T., Methods in Enzymology (Protein Kinase Classification) p. 3, Hunter, T.; Sefton, B. M.; eds. vol. 200, Academic Press; San Diego, 1991].

For most biological responses, multiple intracellular kinases are involved and an individual kinase can be involved in more than one signaling event. These kinases are often cytosolic and can translocate to the nucleus or the ribosomes where they can affect transcriptional and translational events, respectively. The involvement of kinases in transcriptional control is presently much better understood than their effect on translation as illustrated by the studies on growth factor induced signal transduction involving MAP/ERK kinase [Marshall, C. J. Cell, **80**, 179 (1995); Herskowitz, I. Cell, **80**, 187 (1995); Hunter, T. Cell, **80**, 225 (1995); Seger, R., and Krebs, E. G. FASEB J., **9**, 726-735 (1995)].

While many signaling pathways are part of cell homeostasis, numerous cytokines (e.g., IL-1 and TNF) and certain other mediators of inflammation (e.g., COX-2, and iNOS) are produced only as a response to stress signals such as bacterial lipopolysaccharide (LPS). The first indications suggesting that the signal transduction pathway leading to LPS-induced cytokine biosynthesis involved protein kinases came from studies of Weinstein [Weinstein, *et al.*, J. Immunol. **151**, 3829(1993)] but the specific protein kinases involved were not identified. Working from a similar

perspective, Han [Han, *et al.*, *Science* **265**, 808(1994)] identified murine p38 as a kinase which is tyrosine phosphorylated in response to LPS. Definitive proof of the involvement of the p38 kinase in LPS-stimulated signal transduction pathway leading to the initiation of proinflammatory cytokine biosynthesis was provided by the independent discovery of p38 kinase by Lee [Lee, *et al.*, *Nature*, **372**, 739(1994)] as the molecular target for a novel class of anti-inflammatory agents. The discovery of p38 (termed by Lee as CSBP 1 and 2) provided a mechanism of action of a class of anti-inflammatory compounds for which SK&F 86002 was the prototypic example. These compounds inhibited IL-1 and TNF synthesis in human monocytes at concentrations in the low  $\mu$ M range [Lee, *et al.*, *Int. J. Immunopharmac.* 10(7), 835(1988)] and exhibited activity in animal models which are refractory to cyclooxygenase inhibitors [Lee, *et al.*, *Annals N. Y. Acad. Sci.*, **696**, 149(1993)].

### MITOGEN AND STRESS ACTIVATED PROTEIN KINASE CASCADES




Figure 1

It is now firmly established that CSBP/p38 is one of several kinases involved in a stress-response signal transduction pathway which is parallel to and largely independent of the analogous mitogen-activated protein kinase (MAP) kinase cascade (Figure 1). Stress signals, including LPS, pro-inflammatory cytokines, oxidants, UV light and osmotic stress, activate kinases upstream from CSBP/p38 which in turn

phosphorylate CSBP/p38 at threonine 180 and tyrosine 182 resulting in CSBP/p38 activation. MAPKAP kinase-2 and MAPKAP kinase-3 have been identified as downstream substrates of CSBP/p38 which in turn phosphorylate heat shock protein Hsp 27 (Figure 2). It is not yet known whether MAPKAP-2, MAPKAP-3, Mnk1 or Mnk2 are involved in cytokine biosynthesis or alternatively that inhibitors of CSBP/p38 kinase might regulate cytokine biosynthesis by blocking a yet unidentified substrate downstream from CSBP/p38 [Cohen, P. *Trends Cell Biol.*, 353-361(1997)].

## p38 Kinase Pathway




Figure 2

What is known, however, is that in addition to inhibiting IL-1 and TNF, CSBP/p38 kinase inhibitors (SK&F 86002 and SB 203580) also decrease the synthesis of a wide variety of pro-inflammatory proteins including, IL-6, IL-8, GM-CSF and COX-2. Inhibitors of CSBP/p38 kinase have also been shown to suppress the TNF-induced expression of VCAM-1 on endothelial cells, the TNF-induced phosphorylation and activation of cytosolic PLA2 and the IL-1-stimulated synthesis of collagenase and stromelysin. These and additional data demonstrate that CSBP/p38 is involved not only

cytokine synthesis, but also in cytokine signaling [CSBP/P38 kinase reviewed in Cohen, P. Trends Cell Biol., 353-361(1997)].

Interleukin-1 (IL-1) and Tumor Necrosis Factor (TNF) are biological substances produced by a variety of cells, such as monocytes or macrophages. IL-1 has been demonstrated to mediate a variety of biological activities thought to be important in immunoregulation and other physiological conditions such as inflammation [See, e.g., Dinarello et al., *Rev. Infect. Disease*, **6**, 51 (1984)]. The myriad of known biological activities of IL-1 include the activation of T helper cells, induction of fever, stimulation of prostaglandin or collagenase production, neutrophil chemotaxis, induction of acute phase proteins and the suppression of plasma iron levels.

There are many disease states in which excessive or unregulated IL-1 production is implicated in exacerbating and/or causing the disease. These include rheumatoid arthritis, osteoarthritis, endotoxemia and/or toxic shock syndrome, other acute or chronic inflammatory disease states such as the inflammatory reaction induced by endotoxin or inflammatory bowel disease; tuberculosis, atherosclerosis, muscle degeneration, cachexia, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis, and acute synovitis. Recent evidence also links IL-1 activity to diabetes and pancreatic  $\beta$  cells [review of the biological activities which have been attributed to IL-1 Dinarello, *J. Clinical Immunology*, **5** (5), 287-297 (1985)].

Excessive or unregulated TNF production has been implicated in mediating or exacerbating a number of diseases including rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions; sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoisosis, bone resorption diseases, reperfusion injury, graft vs. host reaction, allograft rejections, fever and myalgias due to infection, such as influenza, cachexia secondary to infection or malignancy, cachexia, secondary to acquired immune deficiency syndrome (AIDS), AIDS, ARC (AIDS related complex), keloid formation, scar tissue formation, Crohn's disease, ulcerative colitis, or pyresis.

Interleukin-8 (IL-8) is a chemotactic factor produced by several cell types including mononuclear cells, fibroblasts, endothelial cells, and keratinocytes. Its production from endothelial cells is induced by IL-1, TNF, or lipopolysachharide (LPS). IL-8 stimulates a number of functions in vitro. It has been shown to have chemoattractant properties for neutrophils, T-lymphocytes, and basophils. In addition it induces histamine release from basophils from both normal and atopic individuals as well as lysozomal enzyme release and respiratory burst from neutrophils. IL-8 has also been shown to increase the surface expression of Mac-1 (CD11b/CD18) on neutrophils

without de novo protein synthesis, this may contribute to increased adhesion of the neutrophils to vascular endothelial cells. Many diseases are characterized by massive neutrophil infiltration. Conditions associated with an increased in IL-8 production (which is responsible for chemotaxis of neutrophil into the inflammatory site) would benefit by compounds which are suppressive of IL-8 production.

IL-1 and TNF affect a wide variety of cells and tissues and these cytokines as well as other leukocyte derived cytokines are important and critical inflammatory mediators of a wide variety of disease states and conditions. The inhibition of these cytokines is of benefit in controlling, reducing and alleviating many of these disease states.

Inhibition of signal transduction via CSBP/p38, which in addition to IL-1, TNF and IL-8 described above is also required for the synthesis and/or action of several additional pro-inflammatory proteins (i.e., IL-6, GM-CSF, COX-2, collagenase and stromelysin), is expected to be a highly effective mechanism for regulating the excessive and destructive activation of the immune system. This expectation is supported by the potent and diverse anti-inflammatory activities described for CSBP/p38 kinase inhibitors [Badger, *et al.*, J. Pharm. Exp. Thera. **279** (3): 1453-1461 (1996); Griswold, *et al.*, Pharmacol. Comm. **7**, 323-229 (1996)].

There remains a need for treatment in this field, for compounds which are cytokine suppressive anti-inflammatory drugs, i.e. compounds which are capable of inhibiting the CSBP/p38/RK kinase.

#### SUMMARY OF THE INVENTION

This invention relates to the novel compounds of Formula (I) and (II) and pharmaceutical compositions comprising a compound of Formula (I) or (II) and a pharmaceutically acceptable diluent or carrier.

This invention relates to a method of treating a CSBP/RK/p38 kinase mediated disease in a mammal in need thereof, which method comprises administering to said mammal an effective amount of a compound of Formula (I) or (II).

This invention also relates to a method of inhibiting cytokines and the treatment of a cytokine mediated disease, in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of Formula (I) or (II).


This invention more specifically relates to a method of inhibiting the production of IL-1 in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or (II).

This invention more specifically relates to a method of inhibiting the production of IL-6 in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or (II).

5 This invention more specifically relates to a method of inhibiting the production of IL-8 in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or (II).

This invention more specifically relates to a method of inhibiting the production of TNF in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or (II).

10 Accordingly, the present invention provides for a compound of the formula :



wherein

R<sub>1</sub> is 4-pyridyl, 4-pyrimidinyl, 4-pyridazinyl, 1,2,4-triazin-5-yl, 4-quinolyl, 6-isoquinolyl, or quinazolin-4-yl ring, which ring is optionally substituted independently one to three times with Y, NHR<sub>a</sub>, optionally substituted C<sub>1-4</sub> alkyl, 15 halogen, hydroxyl, optionally substituted C<sub>1-4</sub> alkoxy, optionally substituted C<sub>1-4</sub> alkylthio, optionally substituted C<sub>1-4</sub> alkylsulfinyl, CH<sub>2</sub>OR<sub>12</sub>, amino, mono and di- C<sub>1-6</sub> alkyl substituted amino, N(R<sub>10</sub>)C(O)R<sub>b</sub>, N(R<sub>10</sub>)S(O)<sub>2</sub>R<sub>d</sub>, or an N-heterocyclyl ring which ring has from 5 to 7 members and optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>15</sub>;

20 Y is X<sub>1</sub>-R<sub>a</sub>;

X<sub>1</sub> is sulfur or oxygen;

R<sub>a</sub> is C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic, heterocyclylC<sub>1-6</sub> alkyl, heteroaryl, or heteroarylC<sub>1-6</sub>alkyl, wherein each of these moieties may be optionally substituted;

25 R<sub>b</sub> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl;

R<sub>d</sub> is C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl;

30 R<sub>4</sub> is phenyl, naphth-1-yl or naphth-2-yl, or heteroaryl, which is optionally substituted by one to three substituents, each of which is independently selected, and which, for a 4-phenyl, 4-naphth-1-yl, 5-naphth-2-yl or 6-naphth-2-yl substituent, is halogen, cyano, nitro, C(Z)NR<sub>7</sub>R<sub>17</sub>, C(Z)OR<sub>16</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>COR<sub>12</sub>, SR<sub>5</sub>, S(O)R<sub>5</sub>, OR<sub>12</sub>, halo-substituted-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl, ZC(Z)R<sub>12</sub>, NR<sub>10</sub>C(Z)R<sub>16</sub>, or



- (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>10</sub>R<sub>20</sub> and which, for other positions of substitution, is halogen, cyano, nitro, phenyl, -C(Z)NR<sub>13</sub>R<sub>14</sub>, C(Z)OR<sub>3</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>m</sub>"COR<sub>3</sub>, S(O)<sub>m</sub>R<sub>3</sub>, OR<sub>3</sub>, halo-substituted-C<sub>1-4</sub> alkyl, C<sub>1-10</sub> alkyl, ZC(Z)R<sub>3</sub>, optionally substituted phenyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>m</sub>"NR<sub>10</sub>C(Z)R<sub>3</sub>, NR<sub>10</sub>S(O)<sub>m</sub>'R<sub>8</sub>, NR<sub>10</sub>S(O)<sub>m</sub>'NR<sub>7</sub>R<sub>17</sub>, or
- 5 (CR<sub>10</sub>R<sub>20</sub>)<sub>m</sub>"NR<sub>13</sub>R<sub>14</sub>;
- n is 0, or an integer having a value of 1 to 10;
- v is 0, or an integer having a value of 1 or 2;
- m is 0, or the integer 1 or 2;
- m' is an integer having a value of 1 or 2,
- 10 m" is 0, or an integer having a value of 1 to 5;
- R<sub>2</sub> is hydrogen, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub> OR<sub>9</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OR<sub>11</sub>, C<sub>1-10</sub>alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub>alkyl, heterocyclyl, heterocyclylC<sub>1-10</sub> alkyl,
- 15 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NO<sub>2</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>S(O)<sub>m</sub>'NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OC(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)OR<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)NR<sub>11</sub>OR<sub>9</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>,
- 20 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>N(OR<sub>6</sub>)C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>N(OR<sub>6</sub>)C(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(=NOR<sub>6</sub>)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>19</sub>)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OC(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>10</sub>, 5-(R<sub>18</sub>)-1,2,4-oxadiazol-3-yl or 4-(R<sub>12</sub>)-5-(R<sub>18</sub>R<sub>19</sub>)-4,5-dihydro-1,2,4-oxadiazol-3-yl; wherein the cycloalkyl, cycloalkyl alkyl, aryl, arylalkyl, heteroaryl, heteroaryl alkyl, heterocyclic and heterocyclic alkyl groups may be optionally substituted;
- R<sub>3</sub> is heterocyclyl, heterocyclylC<sub>1-10</sub> alkyl or R<sub>8</sub>;
- R<sub>5</sub> is hydrogen, C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl or NR<sub>7</sub>R<sub>17</sub>, excluding the moieties SR<sub>5</sub> being SNR<sub>7</sub>R<sub>17</sub> and SOR<sub>5</sub> being SOH;
- 30 R<sub>6</sub> is hydrogen, a pharmaceutically acceptable cation, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclic, aroyl, or C<sub>1-10</sub> alkanoyl;
- R<sub>7</sub> and R<sub>17</sub> is each independently selected from hydrogen or C<sub>1-4</sub> alkyl or R<sub>7</sub> and R<sub>17</sub> together with the nitrogen to which they are attached form a heterocyclic ring of 5
- 35 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>15</sub>;

- R<sub>8</sub> is C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>11</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>13</sub>R<sub>14</sub>; wherein the aryl, arylalkyl, heteroaryl, heteroaryl alkyl may  
 5 be optionally substituted;  
 R<sub>9</sub> is hydrogen, C(Z)R<sub>11</sub> or optionally substituted C<sub>1-10</sub> alkyl, S(O)<sub>2</sub>R<sub>18</sub>, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl;  
 R<sub>10</sub> and R<sub>20</sub> is each independently selected from hydrogen or C<sub>1-4</sub> alkyl;  
 R<sub>11</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl,  
 10 aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or heteroarylC<sub>1-10</sub> alkyl, wherein the aryl, arylalkyl, heteroaryl, heteroaryl alkyl, heterocyclyl or heterocyclylalkyl, may be optionally substituted;  
 R<sub>12</sub> is hydrogen or R<sub>16</sub>;  
 R<sub>13</sub> and R<sub>14</sub> is each independently selected from hydrogen or optionally substituted  
 15 C<sub>1-4</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl, or together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub> ;  
 R<sub>15</sub> is hydrogen, C<sub>1-4</sub> alkyl or C(Z)-C<sub>1-4</sub> alkyl;  
 20 R<sub>16</sub> is C<sub>1-4</sub> alkyl, halo-substituted-C<sub>1-4</sub> alkyl, or C<sub>3-7</sub> cycloalkyl;  
 R<sub>18</sub> is C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, aryl, arylC<sub>1-10</sub> alkyl, heterocyclyl, heterocyclyl-C<sub>1-10</sub>alkyl, heteroaryl or heteroarylalkyl, wherein the aryl, arylalkyl, heteroaryl, heteroaryl alkyl, heterocyclyl or heterocyclylalkyl may be optionally substituted;  
 25 R<sub>19</sub> is hydrogen, cyano, C<sub>1-4</sub> alkyl, C<sub>3-7</sub> cycloalkyl or aryl;  
 R<sub>23</sub> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl moiety, all of which may be optionally substituted;  
 Z is oxygen or sulfur;  
 30 provided that when R<sub>1</sub> is a 4-pyridyl, R<sub>4</sub> is a 4-NH<sub>2</sub>S(O)<sub>2</sub>phenyl, then R<sub>2</sub> is other than a 3-position trifluoromethyl group;  
 or a pharmaceutically acceptable salt thereof.

### DETAILED DESCRIPTION OF THE INVENTION

- 35 The novel compounds of Formula (I) or (II) may also be used in association with the veterinary treatment of mammals, other than humans, in need of inhibition of cytokine inhibition or production. In particular, cytokine mediated diseases for

treatment, therapeutically or prophylactically, in animals include disease states such as those noted herein in the Methods of Treatment section, but in particular viral infections. Examples of such viruses include, but are not limited to, lentivirus infections such as, equine infectious anaemia virus, caprine arthritis virus, visna virus, or maedi virus or retrovirus infections, such as but not limited to feline immunodeficiency virus (FIV), bovine immunodeficiency virus, or canine immunodeficiency virus or other retroviral infections.

In Formula (I), suitable R<sub>1</sub> moieties includes 4-pyridyl, 4-pyrimidinyl, 4-pyridazinyl, 1,2,4-triazin-5-yl, 4-quinolyl, 6-isoquinolyl, or 4-quinazolinyl, of which the 4-pyridyl, or 4-pyrimidinyl are preferred. A preferred ring placement of the R<sub>1</sub> substituent on the 4-pyridyl derivative is the 2-position, such as in 2-methoxy-4-pyridyl. A preferred ring placement on the 4-pyrimidinyl ring is also at the 2-position, such as in 2-methoxy-pyrimidinyl.

A preferred embodiment of Formula (I) is where R<sub>1</sub> is an optionally substituted 4-pyridazinyl or 1,2,4-triazin-5-yl ring, substituted as defined above in formula (I).

Suitable additional substituents for the R<sub>1</sub> heteroaryl rings are Y, NHR<sub>a</sub>, optionally substituted C<sub>1-4</sub> alkyl, halogen, hydroxyl, optionally substituted C<sub>1-4</sub> alkoxy, optionally substituted C<sub>1-4</sub> alkylthio, optionally substituted C<sub>1-4</sub> alkylsulfinyl, CH<sub>2</sub>OR<sub>12</sub>, amino, mono and di- C<sub>1-6</sub> alkyl substituted amino, N(R<sub>10</sub>)C(O)R<sub>b</sub>; N(R<sub>10</sub>)S(O)<sub>2</sub>R<sub>d</sub>; or an N-heterocyclyl ring which ring has from 5 to 7 members and optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>15</sub>.

Suitably Y is X<sub>1</sub>-R<sub>a</sub>; and X<sub>1</sub> is oxygen or sulfur, preferably oxygen.

Suitably R<sub>a</sub> is C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic, heterocyclylC<sub>1-6</sub>alkyl, heteroaryl, or heteroarylC<sub>1-6</sub>alkyl, wherein each of these moieties may be optionally substituted as defined herein.

When R<sub>a</sub> is aryl, it is preferably phenyl or naphthyl. When R<sub>a</sub> is an arylC<sub>1-6</sub>alkyl, it is preferably benzyl or naphthylmethyl. When R<sub>a</sub> is a heterocyclic or heterocyclylC<sub>1-6</sub>alkyl moiety, the heterocyclic portion is preferably pyrrolidinyl, piperazine, piperidine, morpholino, tetrahydropyran, tetrahydrothiopyran, tetrahydrothiopyran-sulfinyl, tetrahydrothio-pyransulfonyl, pyrrolidinyl, indole, or piperonyl. It is noted that the heterocyclic rings herein may contain unsaturation, such as in an indole ring. When R<sub>a</sub> is a heteroaryl or heteroarylalkyl moiety it is as defined herein in the definition section.

The R<sub>a</sub> aryl, heterocyclic and heteroaryl containing rings may also be optionally substituted one or more times independently, preferably 1 to 3 times, with halogen, such as fluoro or chloro; C<sub>1-4</sub> alkyl, such as methyl, ethyl, propyl, isopropyl, or t-butyl; halosubstituted alkyl, such as CF<sub>3</sub>; hydroxy; hydroxy substituted C<sub>1-4</sub> alkyl; C<sub>1-4</sub>

alkoxy, such as methoxy or ethoxy; S(O)<sub>m</sub>alkyl and S(O)<sub>m</sub> aryl; C(O)OR<sub>11</sub>, such as C(O)C<sub>1-4</sub> alkyl or C(O)OH moieties; C(O)R<sub>11</sub>; OC(O)R<sub>c</sub>; O-(CH<sub>2</sub>)<sub>s</sub>-O-, such as in a ketal or dioxyalkylene bridge, and s is 1 to 3; amino; NR<sub>13</sub>N<sub>14</sub>; N(R<sub>10</sub>)C(O)R<sub>b</sub>; N(R<sub>10</sub>)S(O)<sub>2</sub>R<sub>d</sub>; C(O)NR<sub>10</sub>R<sub>20</sub>; S(O)<sub>2</sub>(CR<sub>10</sub>R<sub>20</sub>)<sub>t</sub>NR<sub>13</sub>R<sub>14</sub> (wherein t is 0, or an integer of 1 to 3); cyano; nitro; aryl, such as phenyl; an optionally substituted arylC<sub>1-6</sub>alkyl, such as benzyl or phenethyl; aryloxy, such as phenoxy; or arylC<sub>1-6</sub>alkyloxy such as benzyloxy.

Suitably, R<sub>b</sub> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl, wherein all of these moieties may be optionally substituted.

Suitably, R<sub>c</sub> is hydrogen, optionally substituted C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl moiety, wherein all of these moieties may be optionally substituted.

Suitably, R<sub>d</sub> is C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl, wherein all of these moieties may be optionally substituted.

When the R<sub>a</sub> moiety is an alkyl group it may be optionally substituted as defined herein in the definition section below. Also, the alkyl portion of the R<sub>1</sub> substituents, where applicable, such as the mono- and di-C<sub>1-6</sub> alkyl amino moieties, may also be halo substituted.

Preferably, the R<sub>a</sub> group is an alkyl, such as methyl, an optionally substituted aryl, such as phenyl, or an optionally substituted arylalkyl, such as benzyl.

When the R<sub>1</sub> optional substituent group is N(R<sub>10</sub>)C(O) R<sub>b</sub>, R<sub>b</sub> is preferably a C<sub>1-6</sub> alkyl; and R<sub>10</sub> is preferably hydrogen. It is also recognized that all the R<sub>b</sub> moieties, in particular the C<sub>1-6</sub> alkyl group may be optionally substituted, preferably from one to three times as defined herein. Preferably R<sub>b</sub> is C<sub>1-6</sub> alkyl substituted with halogen, such as fluorine, as in trifluoromethyl or trifluoroethyl.

Suitably, R<sub>4</sub> is phenyl, naphth-1-yl or naphth-2-yl, a heteroaryl ring or a fused phenyl containing ring system. Preferably, R<sub>4</sub> is a phenyl or naphthyl ring.

Suitable substitutions for R<sub>4</sub> when this is a 4-phenyl, 4-naphth-1-yl, 5-naphth-2-yl or 6-naphth-2-yl moiety are one to three, preferably one to two, substituents each of which are independently selected from halogen, SR<sub>5</sub>, SOR<sub>5</sub>, OR<sub>12</sub>, CF<sub>3</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>10</sub>R<sub>20</sub>, and for other positions of substitution on these rings preferred

substitution is halogen,  $S(O)_mR_3$ ,  $OR_3$ ,  $CF_3$ ,  $(CR_{10}R_{20})_mNR_{13}R_{14}$ ,  $NR_{10}C(Z)R_3$  and  $NR_{10}S(O)_mR_8$ . Preferred substituents for the 4-position in phenyl and naphth-1-yl and on the 5-position in naphth-2-yl include halogen, especially fluoro and chloro, and  $SR_5$  and  $SOR_5$  wherein  $R_5$  is preferably a  $C_{1-2}$  alkyl, more preferably methyl; of which the fluoro and chloro is more preferred, and most especially preferred is fluoro. For all other substituents, including the heteroaryl ring(s) and fused phenyl ring systems, or the 3-position in phenyl and naphth-1-yl rings, the substituents are independently selected from halogen, especially fluoro and chloro;  $OR_3$ , especially  $C_{1-4}$  alkoxy;  $CF_3$ ,  $NR_{10}R_{20}$ , such as amino;  $NR_{10}C(Z)R_3$ , especially  $NHCO(C_{1-10}alkyl)$ ;  $NR_{10}S(O)_mR_8$ , especially  $NHSO_2(C_{1-10}alkyl)$ ; and  $SR_3$  and  $SOR_3$  wherein  $R_3$  is preferably a  $C_{1-2}$  alkyl, more preferably methyl.

When the phenyl ring is disubstituted, preferably it is two independent halogen moieties, such as fluoro and chloro, preferably di-chloro and more preferably in the 3, 4-position. It is also preferred that for the 3-position of both the  $OR_3$  and  $ZC(Z)R_3$  moieties, that the  $R_3$  may also include hydrogen.

Preferably, the  $R_4$  moiety is an unsubstituted or substituted phenyl moiety. More preferably,  $R_4$  is phenyl or phenyl substituted at the 4-position with fluoro and/or substituted at the 3-position with fluoro, chloro,  $C_{1-4}$  alkoxy, methanesulfonamido or acetamido, or  $R_4$  is a phenyl di-substituted at the 3,4-position independently with chloro or fluoro, more preferably chloro. Most preferably,  $R_4$  is 4-fluorophenyl. Preferably  $R_4$  is not substituted with  $S(O)_2R_3$  when  $R_3$  is  $NR_{13}NR_{14}$ , and  $R_{13}$  and  $R_{14}$  are hydrogen, yielding the moiety  $NH_2S(O)_2-$  in the 4-position.

The  $R_4$  substituent may also include a fused phenyl containing ring system, i.e. a benzene ring fused to a second 4-7 membered carboxylic or heterocyclic ring (containing one or more of O/S/N) which additional ring may not be aromatic, i.e. it can contain some unsaturation. This fused ring system can include such rings as a tetrahydronaphthalene ring, an indoline ring, an indole or a benzimidazole ring system.

In Formula (I), Z is suitably oxygen or sulfur.

Suitably,  $R_2$  is hydrogen,  $(CR_{10}R_{23})_nOR_9$ ,  $(CR_{10}R_{23})_nOR_{11}$ ,  $C_{1-10}alkyl$ , halo-substituted  $C_{1-10}alkyl$ ,  $C_{2-10}alkenyl$ ,  $C_{2-10}alkynyl$ ,  $C_{3-7}cycloalkyl$ ,  $C_{3-7}cycloalkylC_{1-10}alkyl$ ,  $C_{5-7}cycloalkenyl$ ,  $C_{5-7}cycloalkenylC_{1-10}alkyl$ , aryl, aryl $C_{1-10}alkyl$ , heteroaryl, heteroaryl $C_{1-10}alkyl$ , heterocyclyl, heterocyclyl $C_{1-10}alkyl$ ,  $(CR_{10}R_{23})_nS(O)_mR_{18}$ ,  $(CR_{10}R_{23})_nNHS(O)_2R_{18}$ ,  $(CR_{10}R_{23})_nNR_{13}R_{14}$ ,  $(CR_{10}R_{23})_nNO_2$ ,  $(CR_{10}R_{23})_nCN$ ,  $(CR_{10}R_{23})_nS(O)_mNR_{13}R_{14}$ ,  $(CR_{10}R_{23})_nC(Z)R_{11}$ ,  $(CR_{10}R_{23})_nOC(Z)R_{11}$ ,  $(CR_{10}R_{23})_nC(Z)OR_{11}$ ,

(CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)NR<sub>11</sub>OR<sub>9</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>11</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>N(OR<sub>6</sub>)C(Z)NR<sub>13</sub>R<sub>14</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>N(OR<sub>6</sub>)C(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(=NOR<sub>6</sub>)R<sub>11</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>19</sub>)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OC(Z)NR<sub>13</sub>R<sub>14</sub>,  
 5 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>10</sub>, 5-(R<sub>18</sub>)-1,2,4-oxadiazol-3-yl or 4-(R<sub>12</sub>)-5-(R<sub>18</sub>R<sub>19</sub>)-4,5-dihydro-1,2,4-oxadiazol-3-yl; wherein the cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl, heteroaryl alkyl, heterocyclic and heterocyclic alkyl groups may be optionally substituted.

10 Suitably, R<sub>23</sub> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl moiety, all of which may be optionally substituted as defined below.

Preferably, R<sub>2</sub> is hydrogen, an optionally substituted heterocyclyl ring, and  
 15 optionally substituted heterocyclylC<sub>1-10</sub> alkyl, an optionally substituted C<sub>1-10</sub> alkyl, an optionally substituted C<sub>3-7</sub>cycloalkyl, an optionally substituted C<sub>3-7</sub>cycloalkyl C<sub>1-10</sub> alkyl, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)OR<sub>11</sub> group, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>13</sub>R<sub>14</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>18</sub>, an optionally substituted aryl; an optionally substituted arylC<sub>1-10</sub> alkyl, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OR<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)R<sub>11</sub>, or  
 20 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(=NOR<sub>6</sub>)R<sub>11</sub> group.

Preferably, R<sub>2</sub> is selected from hydrogen, C<sub>1-10</sub> alkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylC<sub>1-10</sub> alkyl, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NS(O)<sub>2</sub>R<sub>18</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>18</sub>, arylC<sub>1-10</sub> alkyl, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>13</sub>R<sub>14</sub>, optionally substituted  
 25 C<sub>3-7</sub>cycloalkyl, or optionally substituted C<sub>3-7</sub>cycloalkyl C<sub>1-10</sub> alkyl.

For compounds of Formula (I) only, when R<sub>2</sub> is substituted in the 3-position, than R<sub>2</sub> is other than a halosubstituted alkyl, such as CF<sub>3</sub>. Suitably, when R<sub>1</sub> is a 4-pyridyl, and R<sub>4</sub> is a 4-NH<sub>2</sub>S(O)<sub>2</sub>phenyl, then R<sub>2</sub> is other than a 3-position  
 30 trifluoromethyl group;

When R<sub>2</sub> is an optionally substituted heterocyclyl, the ring is preferably a morpholino, pyrrolidinyl, piperazinyl, or a piperidinyl group. When the ring is optionally substituted, the substituents may be directly attached to the free nitrogen,  
 35 such as in the piperidinyl group or pyrrole ring, or on the ring itself. Preferably the ring is a piperidine or pyrrole, more preferably piperidine. The heterocyclyl ring may be optionally substituted one to four times independently by halogen; C<sub>1-4</sub> alkyl; aryl,

such as phenyl; aryl alkyl, such as benzyl - wherein the aryl or aryl alkyl moieties themselves may be optionally substituted (as in the definition section below); C(O)OR<sub>11</sub>, such as the C(O)C<sub>1-4</sub> alkyl or C(O)OH moieties; C(O)H; C(O)C<sub>1-4</sub> alkyl, hydroxy substituted C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, S(O)<sub>m</sub>C<sub>1-4</sub> alkyl, or NR<sub>10</sub>R<sub>20</sub>.

5

Preferably if the ring is a piperidine, the substituents are directly attached on the available nitrogen, i.e. a 1-Formyl-4-piperidine, 1-benzyl-4-piperidine, 1-methyl-4-piperidine, 1-ethoxycarbonyl-4-piperidine. If the ring is substituted by an alkyl group and the ring is attached in the 4-position, it is preferably substituted in the 2- or 6-  
10 position or both, such as 2,2,6,6-tetramethyl-4-piperidine.

When R<sub>2</sub> is an optionally substituted heterocyclyl C<sub>1-10</sub> alkyl group, the ring is preferably a morpholino, pyrrolidinyl, piperazinyl or a piperidinyl group. Preferably the alkyl chain is 1 to 4 carbons, more preferably 3 or 4, and most preferably 3, such as  
15 in a propyl group. Preferred heterocyclic alkyl groups include but are not limited to, morpholino ethyl, morpholino propyl, pyrrolidinyl propyl, and piperidinyl propyl moieties.

When R<sub>2</sub> is an optionally substituted C<sub>3-7</sub>cycloalkyl, or an optionally  
20 substituted C<sub>3-7</sub>cycloalkyl C<sub>1-10</sub> alkyl, the cycloalkyl group is preferably a C<sub>3</sub> or C<sub>6</sub> ring, most preferably a C<sub>6</sub> ring, which rings may be optionally substituted. The cycloalkyl rings may be optionally substituted one to three times independently by halogen, such as fluorine, chlorine, bromine or iodine; hydroxy; OC(O)R<sub>b</sub>, C<sub>1-10</sub>  
25 alkoxy, such as methoxy or ethoxy; S(O)<sub>m</sub> alkyl, such as methylthio, methylsulfinyl or methylsulfonyl; S(O)<sub>m</sub>aryl; cyano, nitro; NR<sub>7</sub>R<sub>17</sub>; N(R<sub>10</sub>)C(O)X<sub>1</sub> and X<sub>1</sub> is C<sub>1-4</sub> alkyl, aryl or arylC<sub>1-4</sub>alkyl; C<sub>1-10</sub> alkyl, such as methyl, ethyl, propyl, isopropyl, or t-butyl; optionally substituted alkyl wherein the substituents are halogen, (such as CF<sub>3</sub>), hydroxy, nitro, cyano, amino, NR<sub>7</sub>R<sub>17</sub>, S(O)<sub>m</sub> alkyl and S(O)<sub>m</sub> aryl; optionally  
30 substituted alkylene, such as ethylene or propylene; optionally substituted alkyne, such as ethyne; C(O)OR<sub>11</sub>, such as the free acid or methyl ester derivative; the group R<sub>e</sub>; C(O)H; =O; =N-OR<sub>11</sub>; N(H)-OH (or substituted alkyl or aryl derivatives thereof on the nitrogen or the oxime moiety); N(OR<sub>f</sub>)-C(O)-R<sub>21</sub>; an optionally substituted aryl, such as phenyl; an optionally substituted arylC<sub>1-4</sub>alkyl, such as benzyl or phenethyl; an  
35 optionally substituted heterocyclyl or heterocyclic C<sub>1-4</sub>alkyl, and further wherein these aryl, arylalkyl, heterocyclic, and heterocyclic alkyl containing moieties are also optionally substituted one to two times by halogen, hydroxy, C<sub>1-10</sub> alkoxy, S(O)<sub>m</sub>

alkyl, cyano, nitro, amino, mono & di-substituted C<sub>1-6</sub> amino, C<sub>1-10</sub> alkyl, or an halosubstituted C<sub>1-10</sub> alkyl.

Suitably R<sub>e</sub> is a 1,3-dioxyalkylene group of the formula -O-(CH<sub>2</sub>)<sub>s</sub>-O-, wherein s is 1 to 3, preferably s is 2 yielding a 1,3-dioxyethylene moiety, or ketal functionality.

5 Suitably R<sub>f</sub> is hydrogen, a pharmaceutically acceptable cation, aroyl or a C<sub>1-10</sub> alkanoyl group.


Suitably R<sub>21</sub> is NR<sub>22</sub>R<sub>24</sub>; alkyl 1-6; halosubstituted alkyl 1-6; hydroxy substituted alkyl 1-6; alkenyl 2-6; aryl or heteroaryl optionally substituted by halogen, alkyl 1-6, halosubstituted alkyl 1-6, hydroxyl, or alkoxy 1-6.

10 Suitably R<sub>22</sub> is H or alkyl 1-6.

Suitably R<sub>24</sub> is H, alkyl 1-6, aryl, benzyl, heteroaryl, alkyl substituted by halogen or hydroxyl, or phenyl substituted by a member selected from the group consisting of halo, cyano, alkyl 1-12, alkoxy 1-6, halosubstituted alkyl 1-6, alkylthio, alkylsulphonyl, or alkylsulfinyl; or R<sub>22</sub> and R<sub>24</sub> may together with the nitrogen to  
15 which they are attached form a ring having 5 to 7 members, which ring members may be optionally replaced by a heteroatom selected from oxygen, sulfur or nitrogen. The ring may be saturated or contain more than one unsaturated bond. Preferably R<sub>21</sub> is NR<sub>22</sub>R<sub>24</sub> and R<sub>22</sub> and R<sub>24</sub> are preferably hydrogen.

20 When the R<sub>2</sub> cycloalkyl moiety is substituted by NR<sub>7</sub>R<sub>17</sub> group, or NR<sub>7</sub>R<sub>17</sub> C<sub>1-10</sub> alkyl group, and the R<sub>7</sub> and R<sub>17</sub> are as defined in Formula (I), the substituent is preferably an amino, amino alkyl, or an optionally substituted pyrrolidinyl moiety. In those cases where NR<sub>7</sub>R<sub>17</sub> and NR<sub>13</sub>R<sub>14</sub> together cyclize to form a 5 to 7 membered ring, it is noted that those rings may be optionally substituted 1 to 3 times as defined in  
25 the definition section.

A preferred position of ring substitution on the C<sub>6</sub> cycloalkyl moiety is the 4-position. When the cycloalkyl ring is di-substituted it is preferably di-substituted at the 4- position, such as in:



30

wherein R<sup>1'</sup> and R<sup>2'</sup> are independently the optional substituents indicated above for R<sub>2</sub>.



Preferably, R<sup>1'</sup> and R<sup>2'</sup> are hydrogen, hydroxy, alkyl, substituted alkyl, optionally substituted alkyne, aryl, arylalkyl, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>7</sub>R<sub>17</sub>, and N(R<sub>10</sub>)C(O)R<sub>11</sub>.

Suitably, alkyl is C<sub>1-4</sub> alkyl, such as methyl, ethyl, or isopropyl; NR<sub>7</sub>R<sub>17</sub> and NR<sub>7</sub>R<sub>17</sub> alkyl, such as amino, methylamino, aminomethyl, aminoethyl; substituted alkyl such as  
5 in cyanomethyl, cyanoethyl, nitroethyl; pyrrolidinyl; aryl such as in phenyl; arylalkyl, such as in benzyl; optionally substituted alkyne, such as ethyne or propynyl; or together R<sup>1'</sup> and R<sup>2'</sup> are a keto functionality.

In all instances herein where there is an alkenyl or alkynyl moiety as a  
10 substituent group, the unsaturated linkage, i.e., the vinylene or acetylene linkage is preferably not directly attached to the nitrogen, oxygen or sulfur moieties, for instance in OR<sub>3</sub>, or for certain R<sub>2</sub> moieties.

Suitable pharmaceutically acceptable salts are well known to those skilled in the  
15 art and include basic salts of inorganic and organic acids, such as hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, methane sulphonic acid, ethane sulphonic acid, acetic acid, malic acid, tartaric acid, citric acid, lactic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid and mandelic acid. In addition, pharmaceutically acceptable salts of compounds of Formula  
20 (I) may also be formed with a pharmaceutically acceptable cation, for instance, if a substituent group comprises a carboxy moiety. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations.

As used herein, "optionally substituted", unless specifically defined, shall mean  
25 such groups as halogen, such as fluorine, chlorine, bromine or iodine; hydroxy; hydroxy substituted C<sub>1-10</sub>alkyl; C<sub>1-10</sub> alkoxy, such as methoxy or ethoxy; S(O)<sub>m</sub> alkyl, wherein m is 0, 1 or 2, such as methylthio, methylsulfinyl or methylsulfonyl; halosubstituted C<sub>1-10</sub> alkoxy; NR<sub>7</sub>R<sub>17</sub>; C<sub>1-10</sub> alkyl, cycloalkyl, or cycloalkyl alkyl  
30 group, such as methyl, ethyl, propyl, isopropyl, t-butyl, etc. or cyclopropyl methyl; halosubstituted C<sub>1-10</sub> alkyl, such CF<sub>3</sub>; an optionally substituted aryl, such as phenyl, or an optionally substituted arylalkyl, such as benzyl or phenethyl, and wherein these aryl containing moieties may also be optionally substituted one to three times by halogen; hydroxy; hydroxy substituted alkyl; C<sub>1-10</sub> alkoxy; S(O)<sub>m</sub> alkyl; amino, mono & di-C<sub>1-  
35 6</sub> alkyl substituted amino; C<sub>1-6</sub> alkyl; or CF<sub>3</sub>.

The following terms, as used herein, refer to:

- "halo" or "halogens", include the halogens: chloro, fluoro, bromo and iodo.
- "C<sub>1-10</sub>alkyl" or "alkyl" - both straight and branched chain radicals of 1 to 10 carbon atoms, unless the chain length is otherwise limited, including, but not limited to, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, *n*-pentyl and the like.
- The term "cycloalkyl" is used herein to mean cyclic radicals, preferably of 3 to 8 carbons, including but not limited to cyclopropyl, cyclopentyl, cyclohexyl, and the like.
- The term "cycloalkenyl" is used herein to mean cyclic radicals, preferably of 5 to 8 carbons, which have at least one bond including but not limited to cyclopentenyl, cyclohexenyl, and the like.
- The term "alkenyl" is used herein at all occurrences to mean straight or branched chain radical of 2-10 carbon atoms, unless the chain length is limited thereto, including, but not limited to ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl and the like.
- The term "aryl" is used herein to mean, phenyl and naphthyl.
- The term "heteroaryl" (on its own or in any combination, such as "heteroaryloxy", or "heteroaryl alkyl") is used herein to mean a 5-10 membered aromatic ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O or S, such as, but not limited, to pyrrole, pyrazole, furan, thiophene, indole, quinoline, isoquinoline, quinazolinyl, pyridine, pyrimidine, oxazole, thiazole, thiadiazole, triazole, imidazole, or benzimidazole.
- The term "heterocyclic" (on its own or in any combination, such as "heterocyclalkyl") is used herein to mean a saturated or partially unsaturated 4-10 membered ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O, or S; such as, but not limited to, pyrrolidine, piperidine, piperazine, morpholine, tetrahydropyran, or imidazolidine.
- The term "aralkyl" or "heteroarylalkyl" or "heterocyclalkyl" is used herein to mean C<sub>1-4</sub> alkyl as defined above attached to an aryl, heteroaryl or heterocyclic moiety as also defined herein unless otherwise indicate.
- The term "sulfinyl" is used herein to mean the oxide S (O) of the corresponding sulfide, the term "thio" refers to the sulfide, and the term "sulfonyl" refers to the fully oxidized S(O)<sub>2</sub> moiety.
- The term "aroyl" is used herein to mean a C(O)Ar group, wherein Ar is as phenyl, naphthyl, or aryl C<sub>1-4</sub> alkyl derivative such as defined above, such group include but are not limited to benzyl and phenethyl.


• The term "alkanoyl" is used herein to mean a C(O)C<sub>1-10</sub> alkyl wherein the alkyl is as defined above.

It is recognized that the compounds of the present invention may exist as stereoisomers, regioisomers, or diastereoisomers. These compounds may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. All of these compounds are included within the scope of the present invention.

Exemplified compounds of Formula (I) include:

4-[1-(4-Fluorophenyl)-3-phenyl-1*H*-pyrazol-5-yl]pyridine  
 4-[4-Bromo-1-(4-fluorophenyl)-3-phenyl-1*H*-pyrazol-5-yl]pyridine  
 4-[1-(4-Fluorophenyl)-3-[4-(methylthio)phenyl]-1*H*-pyrazol-5-yl]pyridine  
 4-[1-(4-Fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-1*H*-pyrazol-5-yl]pyridine  
 4-[1-(4-Fluorophenyl)-3-[4-(methylsulfinyl)phenyl]-1*H*-pyrazol-5-yl]pyridine; or a pharmaceutically acceptable salt thereof.

Another aspect of the present invention are the novel compounds of Formula (II) as represented by the structure:



(II)

wherein

R<sub>1</sub> is 4-pyridyl, 4-pyrimidinyl, 4-pyridazinyl, 4-triazin-5-yl, quinolyl, isoquinolinyl, or quinazolin-4-yl ring, which ring is optionally substituted independently one to three times with Y, NHR<sub>a</sub>, optionally substituted C<sub>1-4</sub> alkyl, halogen, hydroxyl, optionally substituted C<sub>1-4</sub> alkoxy, optionally substituted C<sub>1-4</sub> alkylthio, optionally substituted C<sub>1-4</sub> alkylsulfinyl, CH<sub>2</sub>OR<sub>12</sub>, amino, mono and di- C<sub>1-6</sub> alkyl substituted amino, N(R<sub>10</sub>)C(O)R<sub>b</sub>, N(R<sub>10</sub>)S(O)<sub>2</sub>R<sub>d</sub> or an N-heterocyclyl ring which ring has from 5 to 7 members and optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>15</sub>;

Y is X<sub>1</sub>-R<sub>a</sub>;

X<sub>1</sub> is sulfur or oxygen;

R<sub>a</sub> is C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic, heterocyclylC<sub>1-6</sub> alkyl, heteroaryl, or heteroarylC<sub>1-6</sub>alkyl, wherein each of these moieties may be optionally substituted;

- $R_b$  is hydrogen,  $C_{1-6}$  alkyl,  $C_{3-7}$  cycloalkyl, aryl, aryl $C_{1-4}$  alkyl, heteroaryl, heteroaryl $C_{1-4}$ alkyl, heterocyclyl, or heterocyclyl $C_{1-4}$  alkyl;
- $R_d$  is  $C_{1-6}$  alkyl,  $C_{3-7}$  cycloalkyl, aryl, aryl $C_{1-4}$  alkyl, heteroaryl, heteroaryl $C_{1-4}$ alkyl, heterocyclyl, or heterocyclyl $C_{1-4}$  alkyl;
- 5  $R_4$  is phenyl, naphth-1-yl or naphth-2-yl, a fused phenyl containing ring system or a heteroaryl, which is optionally substituted by one to three substituents, each of which is independently selected, and which, for a 4-phenyl, 4-naphth-1-yl, 5-naphth-2-yl or 6-naphth-2-yl substituent, is halogen, cyano, nitro,  $C(Z)NR_7R_{17}$ ,  $C(Z)OR_{16}$ ,  $(CR_{10}R_{20})_vCOR_{12}$ ,  $SR_5$ ,  $SOR_5$ ,  $OR_{12}$ , halo-substituted- $C_{1-4}$  alkyl,
- 10  $C_{1-4}$  alkyl,  $ZC(Z)R_{12}$ ,  $NR_{10}C(Z)R_{16}$ , or  $(CR_{10}R_{20})_vNR_{10}R_{20}$  and which, for other positions of substitution, is halogen, cyano,  $C(Z)NR_{13}R_{14}$ ,  $C(Z)OR_3$ ,  $(CR_{10}R_{20})_mCOR_3$ ,  $S(O)_mR_3$ ,  $OR_3$ , halo-substituted- $C_{1-4}$  alkyl,  $C_{1-4}$  alkyl,  $(CR_{10}R_{20})_mNR_{10}C(Z)R_3$ ,  $NR_{10}S(O)_mR_8$ ,  $NR_{10}S(O)_mNR_7R_{17}$ ,  $ZC(Z)R_3$  or  $(CR_{10}R_{20})_mNR_{13}R_{14}$ ;
- 15  $n$  is 0, or an integer having a value of 1 to 10;  
 $m$  is 0, or the integer 1 or 2;  
 $m'$  is an integer having a value of 1 or 2,  
 $m''$  is 0, or an integer having a value of 1 to 5;  
 $v$  is 0, or an integer having a value of 1 or 2;
- 20  $R_2$  is hydrogen,  $(CR_{10}R_{23})_nOR_9$ ,  $(CR_{10}R_{23})_nOR_{11}$ ,  $C_{1-10}$ alkyl, halo-substituted  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl,  $C_{3-7}$  cycloalkyl,  $C_{3-7}$ cycloalkyl $C_{1-10}$  alkyl,  $C_{5-7}$  cycloalkenyl,  $C_{5-7}$  cycloalkenyl  $C_{1-10}$ alkyl, aryl, aryl $C_{1-10}$  alkyl, heteroaryl, heteroaryl $C_{1-10}$ alkyl, heterocyclyl, heterocyclyl $C_{1-10}$  alkyl,  $(CR_{10}R_{23})_nS(O)_mR_{18}$ ,  $(CR_{10}R_{23})_nNHS(O)_2R_{18}$ ,  $(CR_{10}R_{23})_nNR_{13}R_{14}$ ,
- 25  $(CR_{10}R_{23})_nNO_2$ ,  $(CR_{10}R_{23})_nCN$ ,  $(CR_{10}R_{23})_nS(O)_mNR_{13}R_{14}$ ,  $(CR_{10}R_{23})_nC(Z)R_{11}$ ,  $(CR_{10}R_{23})_nOC(Z)R_{11}$ ,  $(CR_{10}R_{23})_nC(Z)OR_{11}$ ,  $(CR_{10}R_{23})_nC(Z)NR_{13}R_{14}$ ,  $(CR_{10}R_{23})_nC(Z)NR_{11}OR_9$ ,  $(CR_{10}R_{23})_nNR_{10}C(Z)R_{11}$ ,  $(CR_{10}R_{23})_nNR_{10}C(Z)NR_{13}R_{14}$ ,  $(CR_{10}R_{23})_nN(OR_6)C(Z)NR_{13}R_{14}$ ,  $(CR_{10}R_{23})_nN(OR_6)C(Z)R_{11}$ ,
- 30  $(CR_{10}R_{23})_nC(=NOR_6)R_{11}$ ,  $(CR_{10}R_{23})_nNR_{10}C(=NR_{19})NR_{13}R_{14}$ ,  $(CR_{10}R_{23})_nOC(Z)NR_{13}R_{14}$ ,  $(CR_{10}R_{23})_nNR_{10}C(Z)NR_{13}R_{14}$ ,  $(CR_{10}R_{23})_nNR_{10}C(Z)OR_{10}$ , 5-( $R_{18}$ )-1,2,4-oxadiazol-3-yl or 4-( $R_{12}$ )-5-( $R_{18}R_{19}$ )-4,5-dihydro-1,2,4-oxadiazol-3-yl; wherein the cycloalkyl, cycloalkyl alkyl, aryl, arylalkyl, heteroaryl, heteroaryl alkyl, heterocyclic and heterocyclic alkyl groups
- 35 may be optionally substituted;
- $R_3$  is heterocyclyl, heterocyclyl $C_{1-10}$  alkyl or  $R_8$ ;

- R<sub>5</sub> is hydrogen, C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl or NR<sub>7</sub>R<sub>17</sub>, excluding the moieties SR<sub>5</sub> being SNR<sub>7</sub>R<sub>17</sub> and SOR<sub>5</sub> being SOH;
- R<sub>6</sub> is hydrogen, a pharmaceutically acceptable cation, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclic, aroyl, or C<sub>1-10</sub> alkanoyl;
- 5 R<sub>7</sub> and R<sub>17</sub> is each independently selected from hydrogen or C<sub>1-4</sub> alkyl or R<sub>7</sub> and R<sub>17</sub> together with the nitrogen to which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>15</sub>;
- 10 R<sub>8</sub> is C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>11</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>13</sub>R<sub>14</sub>; wherein the aryl, arylalkyl, heteroaryl, heteroaryl alkyl containing moieties may be optionally substituted;
- 15 R<sub>9</sub> is hydrogen, C(Z)R<sub>11</sub> or optionally substituted C<sub>1-10</sub> alkyl, S(O)<sub>2</sub>R<sub>18</sub>, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl;
- R<sub>10</sub> and R<sub>20</sub> is each independently selected from hydrogen or C<sub>1-4</sub> alkyl;
- R<sub>11</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or heteroarylC<sub>1-10</sub> alkyl;
- 20 R<sub>12</sub> is hydrogen or R<sub>16</sub>;
- R<sub>13</sub> and R<sub>14</sub> is each independently selected from hydrogen or optionally substituted C<sub>1-4</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl, or together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from
- 25 oxygen, sulfur or NR<sub>9</sub>;
- R<sub>15</sub> is hydrogen, C<sub>1-4</sub> alkyl or C(Z)-C<sub>1-4</sub> alkyl;
- R<sub>16</sub> is C<sub>1-4</sub> alkyl, halo-substituted-C<sub>1-4</sub> alkyl, or C<sub>3-7</sub> cycloalkyl;
- R<sub>18</sub> is C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, aryl, arylC<sub>1-10</sub> alkyl, heterocyclyl, heterocyclyl-C<sub>1-10</sub>alkyl, heteroaryl or heteroarylalkyl;
- 30 R<sub>19</sub> is hydrogen, cyano, C<sub>1-4</sub> alkyl, C<sub>3-7</sub> cycloalkyl or aryl;
- R<sub>23</sub> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl moiety, all of which may be optionally substituted;
- Z is oxygen or sulfur;
- 35 or a pharmaceutically acceptable salt thereof.

In Formula (II), suitable  $R_1$ ,  $R_2$ , and  $R_4$  variables, etc. for use in Formula (II) compounds are the same as those indicated above for Formula (I) compounds and are defined therein. As will be noted the difference between Formula (I) and Formula (II) compounds lay in the saturation of the core ring system.

5

Exemplified compounds of Formula (II) include:

4-[1-(4-Fluorophenyl)-4,5-dihydro-3-phenyl-1*H*-pyrazol-5-yl]pyridine

4-[1-(4-Fluorophenyl)-4,5-dihydro-3-[4-(methylthio)phenyl]-1*H*-pyrazol-5-yl]pyridine


## 10 Synthetic Methods

The compounds of Formula (I) and (II) may be obtained by applying synthetic procedures, some of which are illustrated in Schemes I and II below. The synthesis provided for in these Schemes is applicable for producing compounds of Formula (I) and (II) having a variety of different  $R_1$ ,  $R_2$ , and  $R_4$  groups which are reacted,

15 employing optional substituents which are suitably protected, to achieve compatibility with the reactions outlined herein. Subsequent deprotection, in those cases, then affords compounds of the nature generally disclosed. Once the pyrazole or pyrazoline nucleus has been established, further compounds of Formula (I) or (II) may be prepared by applying standard techniques for functional group interconversion, well known in the art.


20 For instance:  $C(O)NR_{13}R_{14}$  from  $CO_2CH_3$  by heating with or without catalytic metal cyanide, e.g. NaCN, and  $HNR_{13}R_{14}$  in  $CH_3OH$ ;  $OC(O)R_3$  from OH with e.g.,  $ClC(O)R_3$  in pyridine;  $NR_{10}-C(S)NR_{13}R_{14}$  from  $NHR_{10}$  with an alkylisothiocyanate or thiocyanic acid;  $NR_6C(O)OR_6$  from  $NHR_6$  with the alkyl  
25 chloroformate;  $NR_{10}C(O)NR_{13}R_{14}$  from  $NHR_{10}$  by treatment with an isocyanate, e.g.  $HN=C=O$  or  $R_{10}N=C=O$ ;  $-NR_{10}-C(O)R_8$  from  $NHR_{10}$  by treatment with  $Cl-C(O)R_3$  in pyridine;  $C(=NR_{10})NR_{13}R_{14}$  from  $C(NR_{13}R_{14})SR_3$  with  $H_3NR_3^+OAc^-$  by heating in alcohol;  $C(NR_{13}R_{14})SR_3$  from  $C(S)NR_{13}R_{14}$  with  $R_6-I$  in an inert solvent, e.g. acetone;  $C(S)NR_{13}R_{14}$  (where  $R_{13}$  or  $R_{14}$  is not hydrogen) from  $C(S)NH_2$  with  
30  $HNR_{13}R_{14}-C(=NCN)-NR_{13}R_{14}$  from  $C(=NR_{13}R_{14})-SR_3$  with  $NH_2CN$  by heating in anhydrous alcohol, alternatively from  $C(=NH)-NR_{13}R_{14}$  by treatment with  $BrCN$  and  $NaOEt$  in  $EtOH$ ;  $NR_{10}-C(=NCN)SR_8$  from  $-NHR_{10}$  by treatment with  $(R_8S)_2C=NCN$ ;  $NR_{10}SO_2R_3$  from  $NHR_{10}$  by treatment with  $ClSO_2R_3$  by heating in pyridine;  $NR_{10}C(S)R_3$  from  $NR_{10}C(O)R_8$  by treatment with Lawesson's reagent [2,4-*bis*(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane-2,4-disulfide];  $NR_{10}SO_2CF_3$  from  
35  $NHR_6$  with triflic anhydride and base wherein  $R_3$ ,  $R_6$ ,  $R_{10}$ ,  $R_{13}$  and  $R_{14}$  are as defined in Formula (I) herein.

Compounds of Formula (I) are pyrazole derivatives which may be readily prepared using procedures well known to those of skill in the art and may be prepared by analogous methods to those indicated herein below.



Scheme 1

Scheme I illustrates a regioselective synthetic approach to isomerically pure pyrazoles using  $\text{Co}(\text{OAc})_2$  as the catalyst to effect enone formation. Alternative  
 10 synthetic methods for the synthesis of compound 1-Scheme 1 include preparation of vinyl organostannane or zirconium reagent from the acetylene and coupling with a suitably functionalized heterocycle (for example, 4-bromopyridine) using a Pd catalyst. A variety of additional methodologies are available which insure that a wide variety of substitution can be tolerated in compound 1. Similarly, many substituted aryl  
 15 hydrazines are commercially available or can be readily prepared using standard literature procedures. The pyrazoline, 2, is readily oxidized to 3 with  $\text{MnO}_2$ , but other oxidizing agents may also be used, such as DDQ. The pyrazoline 2 compounds are illustrative of compounds of Formula (II) herein. While the illustration in Scheme I is for the preparation of a particular compound of Formula (I and II) (i.e., Scheme I,  $\text{R}_1 =$   
 20 pyridyl,  $\text{R}_4 = 4\text{-fluorophenyl}$ ,  $\text{R}_2 = \text{phenyl}$ ), generalization of the synthesis to groups claimed as  $\text{R}_1$ ,  $\text{R}_2$ , and  $\text{R}_4$  herein can be achieved by employing the appropriate starting materials.



Scheme II


Scheme II illustrates an alternative synthesis of pyrazoles which in many cases will give regioisomeric mixtures of **II** and **III**. However, by careful adjustment of the conditions it may be possible to alter this ratio to favor the desired product **II**. Oxidation of **II** to the sulfoxide or sulfone activates the heterocyclic ring to nucleophilic substitution. Acceptable nucleophiles are alcohols (compound **V**) phenols, anilines and amines. The nucleophilic substitution occurs under both acidic and neutral conditions in a wide range of solvents (ether, THF, DMF, toluene, glyme, etc.), but is preferably conducted using the alkali metal salt of the nucleophile.

Suitable protecting groups for use in the present invention, are well known in the art and described in many references, for instance, *Protecting Groups in Organic Synthesis*, Greene T W, Wiley-Interscience, New York, 1981.

Pharmaceutically acid addition salts of compounds of formula (I) may be obtained in known manner, for example by treatment thereof with an appropriate amount of acid in the presence of a suitable solvent.



Another aspect of the present invention is a process for producing a compound of Formula (I), which process comprises cyclizing a compound of the formula




wherein  $R_2$  is an optionally substituted phenyl, as defined according to formula (I);

- 5 with a compound of the formula:  $R_4NHNH_2$ , wherein  $R_4$  is as defined for Formula (I), to yield a compound of Formula (I), or if necessary, converting a precursor of  $R_1$ ,  $R_2$  and  $R_4$  to a group  $R_1$ ,  $R_2$  and  $R_4$ .

Another aspect of the present invention is a process for producing a compound of Formula (I), wherein  $R_1$  is an optionally substituted pyridyl, which process

- 10 comprises cyclizing a compound of the formula



wherein  $R_2$  is an optionally substituted phenyl, as defined according to formula (I);

- 15 with a compound of the formula:  $R_4NHNH_2$ , wherein  $R_4$  is as defined for Formula (I), to yield a compound of Formula (I), or if necessary, converting a precursor of  $R_1$ ,  $R_2$  and  $R_4$  to a group  $R_1$ ,  $R_2$  and  $R_4$ .

The invention will now be described by reference to the following examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention.

## 20 Synthetic Examples

### Example 1

#### Phenyl-1-(4-pyridyl)prop-1-en-3-one

- 25 Pyridine (0.6g, 7.5 mmol) and Cobalt (ii) acetate (1.3g, 7.5 mmol) were dissolved in DMF to produce a blue solution of the complex to which was added 4-pyridine carboxaldehyde (5g, 46.7 mmol) and acetophenone (6.2g, 51.4 mmol). The resulting mixture was stirred for 18 hours at 90°C. Most of the DMF was evaporated in vacuo and the residue was purified by flash chromatography (silica gel) with 30% EtoAc/Hexane to afford the titled compound, white solid (5g). ESP+ (Mass Spec) m/z 210 (MH+).

30

**Example 2**1-(4-Fluorophenyl)-3-phenyl-5-(4-pyridyl)-1,2-pyrazoline

Phenyl-1-(4-pyridyl)prop-1-en-3-one (4g, 19.1 mmol) and 4-fluorophenyl hydrazine hydrochloride salt (4.7g, 28.7 mmol) were dissolved in ethanol. Sodium acetate (2.4g, 28.7 mmol) was added. The resulting mixture was stirred for 18 hours at 75°C. Most of the ethanol were evaporated in vacuo. The residue was extracted with EtoAc, washed the organic layer with brine, dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure to yield crude enamine as a brownish solid. (6g). ESP+ (Mass Spec) m/z 318 (MH+). The enamine (4.5g, 14.2 mmol) was dissolved in ethanol and 3N HCl slowly added. The resulting mixture was stirred for 18 hours at 75°C, cooled, poured into water, filtered and the white solid air dried. Recrystallized from EtoAc/Hexane (2:10) to afford the titled compound (4.5g) ESP+ (Mass Spec) m/z 317 (MH+).

**Example 3**1-(4-Fluorophenyl)-3-phenyl-5-(4-pyridyl)pyrazole

1-(4-Fluorophenyl)-3-phenyl-5-(4-pyridyl)-1,2-pyrazoline (0.5g, 1.57 mmol) and Manganase (iv) oxide (0.16g, 1.88 mmol) were dissolved in benzene. The resulting mixture was stirred for 18 hours at 25°C, filtered through celite, and concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel) with 40% EtoAc/Hexane to afford the titled compound, yellow solid (5g). ESP+ (Mass Spec) m/z 316 (MH+).

**Example 4**1-(4-Fluorophenyl)-3-phenyl-4-bromo-5-(4-pyridyl)pyrazole

1-(4-Fluorophenyl)-3-phenyl-5-(4-pyridyl)-1,2-pyrazoline (0.6g, 1.89 mmol) was dissolved in 5% aqueous bromine. The resulting mixture was stirred for 18 hours at 25°C. The residue was neutralized with aqueous NaOH, extracted with EtOAc, washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to produce an off white solid. Recrystallization from EtOAc/Hexane (2:10) afforded the titled compound, (17 mg). ESP+ (Mass Spec) m/z 395 (MH+).

**Example 5**(E)-1-(4-methylthiophenyl)-3-(4-pyridyl)-2-propenone

To a mixture of cobalt (ii) acetate (4g, 23.0 mmol) in pyridine was added 4-pyridine carboxaldehyde (5g, 46.7 mmol) and 4-(methylthio) acetophenone (8g, 48.12 mmol). The resulting mixture was heated for 18 hours at 90°C. Following concentration under

reduced pressure, the residue was purified by flash chromatography (silica gel) with 30% EtoAc/Hexane to afford the titled compound, white solid (3g). ESP+ (Mass Spec) m/z 256 (MH+).

5

### Example 6

#### 1-(4-Fluorophenyl)-3-4-(thiomethylphenyl)-5-(4-pyridyl)-1,2-pyrazoline

(E)-1-(4-methylthiophenyl)-3-(4-pyridyl)-2-propenone (3g, 11.7mmol) and 4-fluorophenylhydrazine hydrochloride salt (3g, 18.5 mmol) were dissolved in ethanol to which sodium acetate (1.5g, 18.5 mmol) was added. The resulting mixture was stirred for 18 hours at 75°C after which the bulk of the ethanol was removed under reduced pressure. The residue was extracted with EtOAc, the organic layer was washed with brine, dried with Na<sub>2</sub>SO<sub>4</sub> and concentrated to yield the crude enamine as a brownish solid (3g). ESP+ (Mass Spec) m/z 363 (MH+). enamine (6g). ESP+ (Mass Spec) m/z 318 (MH+).

The enamine (3g, 8.3 mmol) was dissolved in ethanol and 3N HCl added slowly. The resulting mixture was stirred for 18 hours at 75°C, cooled, and poured into water. The resulting precipitate was filtered and the white solid air dried. Recrystallization from EtoAc/Hexane (2:10) afforded the titled compound (4.5g) ESP+ (Mass Spec) m/z 363 (MH+).

### Example 7

#### 1-(4-Fluorophenyl)-3-4-thiomethylphenyl)-5-(4-pyridyl)pyrazole

1-(4-Fluorophenyl)-3-4-thiomethylphenyl)-5-(4-pyridyl)-1,2-pyrazoline (2.1g, 5.8 mmol) and manganese (iv) oxide (0.66g, 7.6 mmol) were dissolved in benzene. The resulting mixture was stirred for 18 hours at 25°C, filtered through celite, and concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel) with 40% EtoAc/Hexane to afford the titled compound, brownish solid (125 mg). ESP+ (Mass Spec) m/z 362 (MH+).

30

### Example 8

#### 1-(4-Fluorophenyl)-3-(4-methylsulfonylphenyl)-5-(4-pyridyl)pyrazole

1-(4-Fluorophenyl)-3-(4-thiomethylphenyl)-5-(4-pyridyl)pyrazole was dissolved in THF cooled to -10°C and oxone (3.2g, 5.26 mmol) in water (15 ml) was added dropwise (T < 5°C). The resulting mixture was warmed to 20°C over 50 min. and then poured into a vigorously stirred mixture of 10% aqueous NaOH (160 ml), extracted with EtoAc, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to a brownish solid. Recrystallization

from EtoAc/Hexane(1:10) afforded the titled compound (50 mg). ESP+ (Mass Spec) m/z 394 (MH+).

#### METHODS OF TREATMENT

5           The compounds of Formula (I) or (II) or a pharmaceutically acceptable salt thereof can be used in the manufacture of a medicament for the prophylactic or therapeutic treatment of any disease state in a human, or other mammal, which is exacerbated or caused by excessive or unregulated cytokine production by such mammal's cell, such as but not limited to monocytes and/or macrophages.

10           As used herein, unless specifically indicated, compounds of Formula (I) also include compounds of Formula (II).

          Compounds of Formula (I) are capable of inhibiting proinflammatory cytokines, such as IL-1, IL-6, IL-8 and TNF and are therefore of use in therapy. IL-1, IL-6, IL-8 and TNF affect a wide variety of cells and tissues and these cytokines, as well as other  
15   leukocyte-derived cytokines, are important and critical inflammatory mediators of a wide variety of disease states and conditions. The inhibition of these pro-inflammatory cytokines is of benefit in controlling, reducing and alleviating many of these disease states.

20           Accordingly, the present invention provides a method of treating a cytokine-mediated disease which comprises administering an effective cytokine-interfering amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

          In particular, compounds of Formula (I) or a pharmaceutically acceptable salt  
25   thereof are of use in the prophylaxis or therapy of any disease state in a human, or other mammal, which is exacerbated by or caused by excessive or unregulated IL-1, IL-8 or TNF production by such mammal's cell, such as, but not limited to, monocytes and/or macrophages.

          Accordingly, in another aspect, this invention relates to a method of inhibiting  
30   the production of IL-1 in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

          There are many disease states in which excessive or unregulated IL-1  
35   production is implicated in exacerbating and/or causing the disease. These include rheumatoid arthritis, osteoarthritis, stroke, endotoxemia and/or toxic shock syndrome, other acute or chronic inflammatory disease states such as the inflammatory reaction induced by endotoxin or inflammatory bowel disease, tuberculosis, atherosclerosis,

muscle degeneration, multiple sclerosis, cachexia, bone resorption, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis and acute synovitis. Recent evidence also links IL-1 activity to diabetes, pancreatic  $\beta$  cells and Alzheimer's disease.

5 In a further aspect, this invention relates to a method of inhibiting the production of TNF in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

10 Excessive or unregulated TNF production has been implicated in mediating or exacerbating a number of diseases including rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, stroke, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoisosis, bone resorption diseases, such as osteoporosis,  
15 reperfusion injury, graft vs. host reaction, allograft rejections, fever and myalgias due to infection, such as influenza, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency syndrome (AIDS), AIDS, ARC (AIDS related complex), keloid formation, scar tissue formation, inflammatory bowel disease, Crohn's disease, ulcerative colitis and pyresis.

20 Compounds of Formula (I) are also useful in the treatment of viral infections, where such viruses are sensitive to upregulation by TNF or will elicit TNF production *in vivo*. The viruses contemplated for treatment herein are those that produce TNF as a result of infection, or those which are sensitive to inhibition, such as by decreased replication, directly or indirectly, by the TNF inhibiting-compounds of Formula (1).  
25 Such viruses include, but are not limited to HIV-1, HIV-2 and HIV-3, Cytomegalovirus (CMV), Influenza, adenovirus and the Herpes group of viruses, such as but not limited to, Herpes Zoster and Herpes Simplex. Accordingly, in a further aspect, this invention relates to a method of treating a mammal afflicted with a human immunodeficiency virus (HIV) which comprises administering to such mammal an effective TNF  
30 inhibiting amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

Compounds of Formula (I) may also be used in association with the veterinary treatment of mammals, other than in humans, in need of inhibition of TNF production. TNF mediated diseases for treatment, therapeutically or prophylactically, in animals  
35 include disease states such as those noted above, but in particular viral infections. Examples of such viruses include, but are not limited to, lentivirus infections such as, equine infectious anaemia virus, caprine arthritis virus, visna virus, or maedi virus or

retrovirus infections, such as but not limited to feline immunodeficiency virus (FIV), bovine immunodeficiency virus, or canine immunodeficiency virus or other retroviral infections.

5 The compounds of Formula (I) may also be used topically in the treatment or prophylaxis of topical disease states mediated by or exacerbated by excessive cytokine production, such as by IL-1 or TNF respectively, such as inflamed joints, eczema, contact dermatitis, psoriasis and other inflammatory skin conditions such as sunburn; inflammatory eye conditions including conjunctivitis; pyresis, pain and other conditions associated with inflammation.

10 Compounds of Formula (I) have also been shown to inhibit the production of IL-8 (Interleukin-8, NAP). Accordingly, in a further aspect, this invention relates to a method of inhibiting the production of IL-8 in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

15 There are many disease states in which excessive or unregulated IL-8 production is implicated in exacerbating and/or causing the disease. These diseases are characterized by massive neutrophil infiltration such as, psoriasis, inflammatory bowel disease, asthma, cardiac and renal reperfusion injury, adult respiratory distress syndrome, thrombosis and glomerulonephritis. All of these diseases are associated with  
20 increased IL-8 production which is responsible for the chemotaxis of neutrophils into the inflammatory site. In contrast to other inflammatory cytokines (IL-1, TNF, and IL-6), IL-8 has the unique property of promoting neutrophil chemotaxis and activation. Therefore, the inhibition of IL-8 production would lead to a direct reduction in the neutrophil infiltration.

25 The compounds of Formula (I) are administered in an amount sufficient to inhibit cytokine, in particular IL-1, IL-6, IL-8 or TNF, production such that it is regulated down to normal levels, or in some case to subnormal levels, so as to ameliorate or prevent the disease state. Abnormal levels of IL-1, IL-6, IL-8 or TNF, for instance in the context of the present invention, constitute: (i) levels of free (not cell  
30 bound) IL-1, IL-6, IL-8 or TNF greater than or equal to 1 picogram per ml; (ii) any cell associated IL-1, IL-6, IL-8 or TNF; or (iii) the presence of IL-1, IL-6, IL-8 or TNF mRNA above basal levels in cells or tissues in which IL-1, IL-6, IL-8 or TNF, respectively, is produced.

35 The discovery that the compounds of Formula (I) are inhibitors of cytokines, specifically IL-1, IL-6, IL-8 and TNF is based upon the effects of the compounds of Formulas (I) on the production of the IL-1, IL-8 and TNF in *in vitro* assays which are described herein.

As used herein, the term "inhibiting the production of IL-1 (IL-6, IL-8 or TNF)" refers to:

- a) a decrease of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels by inhibition of the *in vivo* release of the cytokine by all cells, including but not limited to monocytes or macrophages;
- b) a down regulation, at the genomic level, of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels;
- c) a down regulation, by inhibition of the direct synthesis of the cytokine (IL-1, IL-6, IL-8 or TNF) as a postranslational event; or
- d) a down regulation, at the translational level, of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels.

As used herein, the term "TNF mediated disease or disease state" refers to any and all disease states in which TNF plays a role, either by production of TNF itself, or by TNF causing another monokine to be released, such as but not limited to IL-1, IL-6 or IL-8. A disease state in which, for instance, IL-1 is a major component, and whose production or action, is exacerbated or secreted in response to TNF, would therefore be considered a disease state mediated by TNF.

As used herein, the term "cytokine" refers to any secreted polypeptide that affects the functions of cells and is a molecule which modulates interactions between cells in the immune, inflammatory or hematopoietic response. A cytokine includes, but is not limited to, monokines and lymphokines, regardless of which cells produce them. For instance, a monokine is generally referred to as being produced and secreted by a mononuclear cell, such as a macrophage and/or monocyte. Many other cells however also produce monokines, such as natural killer cells, fibroblasts, basophils, neutrophils, endothelial cells, brain astrocytes, bone marrow stromal cells, epidermal keratinocytes and B-lymphocytes. Lymphokines are generally referred to as being produced by lymphocyte cells. Examples of cytokines include, but are not limited to, Interleukin-1 (IL-1), Interleukin-6 (IL-6), Interleukin-8 (IL-8), Tumor Necrosis Factor-alpha (TNF- $\alpha$ ) and Tumor Necrosis Factor beta (TNF- $\beta$ ).

As used herein, the term "cytokine interfering" or "cytokine suppressive amount" refers to an effective amount of a compound of Formula (I) which will cause a decrease in the *in vivo* levels of the cytokine to normal or sub-normal levels, when given to a patient for the prophylaxis or treatment of a disease state which is exacerbated by, or caused by, excessive or unregulated cytokine production.

As used herein, the cytokine referred to in the phrase "inhibition of a cytokine, for use in the treatment of a HIV-infected human" is a cytokine which is implicated in (a) the initiation and/or maintenance of T cell activation and/or activated T cell-

mediated HIV gene expression and/or replication and/or (b) any cytokine-mediated disease associated problem such as cachexia or muscle degeneration.

As TNF- $\beta$  (also known as lymphotoxin) has close structural homology with TNF- $\alpha$  (also known as cachectin) and since each induces similar biologic responses and binds to the same cellular receptor, both TNF- $\alpha$  and TNF- $\beta$  are inhibited by the compounds of the present invention and thus are herein referred to collectively as "TNF" unless specifically delineated otherwise.

A new member of the MAP kinase family, alternatively termed CSBP, p38, or RK, has been identified independently by several laboratories recently [See Lee *et al.*, Nature, Vol. 300 n(72), 739-746 (1994)]. Activation of this novel protein kinase via dual phosphorylation has been observed in different cell systems upon stimulation by a wide spectrum of stimuli, such as physicochemical stress and treatment with lipopolysaccharide or proinflammatory cytokines such as interleukin-1 and tumor necrosis factor. The cytokine biosynthesis inhibitors, of the present invention, compounds of Formula (I), have been determined to be potent and selective inhibitors of CSBP/p38/RK kinase activity. These inhibitors are of aid in determining the signaling pathways involvement in inflammatory responses. In particular, for the first time a definitive signal transduction pathway can be prescribed to the action of lipopolysaccharide in cytokine production in macrophages. In addition to those diseases already noted, treatment of stroke, neurotrauma, cardiac and renal reperfusion injury, chronic renal failure, congestive heart failure, cancer, thrombosis, glomerulonephritis, diabetes and pancreatic  $\beta$  cells, multiple sclerosis, muscle degeneration, eczema, psoriasis, sunburn, and conjunctivitis are also included.

The cytokine inhibitors were subsequently tested in a number of animal models for anti-inflammatory activity. Model systems were chosen that were relatively insensitive to cyclooxygenase inhibitors in order to reveal the unique activities of cytokine suppressive agents. The inhibitors exhibited significant activity in many such *in vivo* studies. Most notable are its effectiveness in the collagen-induced arthritis model and inhibition of TNF production in the endotoxic shock model. In the latter study, the reduction in plasma level of TNF correlated with survival and protection from endotoxic shock related mortality. Also of great importance are the compounds effectiveness in inhibiting bone resorption in a rat fetal long bone organ culture system. Griswold *et al.*, (1988) *Arthritis Rheum.* **31**:1406-1412; Badger, *et al.*, (1989) *Circ. Shock* **27**, 51-61; Votta *et al.*, (1994) *in vitro. Bone* **15**, 533-538; Lee *et al.*, (1993), *B Ann. N. Y. Acad. Sci.* **696**, 149-170.



Another aspect of the present invention is to the novel use of these CSBP/cytokine inhibitors for the treatment of chronic inflammatory or proliferative or angiogenic diseases which are caused by excessive, or inappropriate angiogenesis.

Chronic diseases which have an inappropriate angiogenic component are  
5 various ocular neovascularizations, such as diabetic retinopathy and macular degeneration. Other chronic diseases which have an excessive or increased proliferation of vasculature are tumor growth and metastasis, atherosclerosis, and certain arthritic conditions. Therefore cytokine inhibitors will be of utility in the blocking of the angiogenic component of these disease states.

10 The term "excessive or increased proliferation of vasculature inappropriate angiogenesis" as used herein includes, but is not limited to, diseases which are characterized by hemangiomas and ocular diseases.

The term "inappropriate angiogenesis" as used herein includes, but is not limited to, diseases which are characterized by vesicle proliferation with accompanying tissue  
15 proliferation, such as occurs in cancer, metastasis, arthritis and atherosclerosis.

In order to use a compound of Formula (I) or a pharmaceutically acceptable salt thereof in therapy, it will normally be Formulated into a pharmaceutical composition in accordance with standard pharmaceutical practice. This invention, therefore, also  
20 relates to a pharmaceutical composition comprising an effective, non-toxic amount of a compound of Formula (I) and a pharmaceutically acceptable carrier or diluent.

Compounds of Formula (I), pharmaceutically acceptable salts thereof and pharmaceutical compositions incorporating such may conveniently be administered by any of the routes conventionally used for drug administration, for instance, orally,  
25 topically, parenterally or by inhalation. The compounds of Formula (I) may be administered in conventional dosage forms prepared by combining a compound of Formula (I) with standard pharmaceutical carriers according to conventional procedures. The compounds of Formula (I) may also be administered in conventional dosages in combination with a known, second therapeutically active compound. These  
30 procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation. It will be appreciated that the form and character of the pharmaceutically acceptable character or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables. The carrier(s) must be "acceptable" in  
35 the sense of being compatible with the other ingredients of the Formulation and not deleterious to the recipient thereof.

The pharmaceutical carrier employed may be, for example, either a solid or liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are syrup, peanut oil, olive oil, water and the like. Similarly, the carrier or  
5 diluent may include time delay material well known to the art, such as glyceryl mono-stearate or glyceryl distearate alone or with a wax.

A wide variety of pharmaceutical forms can be employed. Thus, if a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier  
10 will vary widely but preferably will be from about 25 mg. to about 1g. When a liquid carrier is used, the preparation will be in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid suspension.

Compounds of Formula (I) may be administered topically, that is by non-systemic administration. This includes the application of a compound of Formula (I)  
15 externally to the epidermis or the buccal cavity and the instillation of such a compound into the ear, eye and nose, such that the compound does not significantly enter the blood stream. In contrast, systemic administration refers to oral, intravenous, intraperitoneal and intramuscular administration.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of inflammation such as  
20 liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose. The active ingredient may comprise, for topical administration, from 0.001% to 10% w/w, for instance from 1% to 2% by weight of the Formulation. It may however comprise as much as 10% w/w but preferably will comprise less than 5%  
25 w/w, more preferably from 0.1% to 1% w/w of the Formulation.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include  
30 an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid Formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely-divided or powdered form, alone or in solution or  
35 suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy base. The base may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin

such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives or a fatty acid such as steric or oleic acid together with an alcohol such as propylene glycol or a macrogel. The Formulation may incorporate any suitable surface active agent such as an anionic, cationic or non-ionic surfactant such as a sorbitan ester or a  
5 polyoxyethylene derivative thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active ingredient in a  
10 suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining at 98-100° C. for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container  
15 by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol and propylene glycol.

Compounds of formula (I) may be administered parenterally, that is by  
20 intravenous, intramuscular, subcutaneous intranasal, intrarectal, intravaginal or intraperitoneal administration. The subcutaneous and intramuscular forms of parenteral administration are generally preferred. Appropriate dosage forms for such administration may be prepared by conventional techniques. Compounds of Formula (I) may also be administered by inhalation, that is by intranasal and oral inhalation  
25 administration. Appropriate dosage forms for such administration, such as an aerosol Formulation or a metered dose inhaler, may be prepared by conventional techniques.

For all methods of use disclosed herein for the compounds of Formula (I), the daily oral dosage regimen will preferably be from about 0.1 to about 80 mg/kg of total body weight, preferably from about 0.2 to 30 mg/kg, more preferably from about 0.5  
30 mg to 15 mg. The daily parenteral dosage regimen about 0.1 to about 80 mg/kg of total body weight, preferably from about 0.2 to about 30 mg/kg, and more preferably from about 0.5 mg to 15 mg/kg. The daily topical dosage regimen will preferably be from 0.1 mg to 150 mg, administered one to four, preferably two or three times daily. The daily inhalation dosage regimen will preferably be from about 0.01 mg/kg to about 1  
35 mg/kg per day. It will also be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of a compound of Formula (I) or a pharmaceutically acceptable salt thereof will be determined by the nature and extent of

the condition being treated, the form, route and site of administration, and the particular patient being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of a compound of Formula (I) or a  
5 pharmaceutically acceptable salt thereof given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

The invention will now be described by reference to the following biological examples which are merely illustrative and are not to be construed as a limitation of the  
10 scope of the present invention.

### BIOLOGICAL EXAMPLES

The cytokine-inhibiting effects of compounds of the present invention were determined by the following *in vitro* assays:

15 Interleukin - 1 (IL-1), Interleukin -8 (IL-8 ), and Tumour Necrosis Factor (TNF) assays may be found in a number of publications, in particular suitable assays for use herein are described in Adams et al., US 5,593,992, whose disclosure is incorporated by reference.

#### 20 **In vivo TNF assay:**

While the above indicated assay in an *in vitro* assay, the compounds of Formula (I) may also be tested in an *in vivo* system such as described in :

(1) Griswold *et al.*, Drugs Under Exp. and Clinical Res., XIX (6), 243-48 (1993);

or

25 (2) Boehm, *et al.*, *Journal Of Medicinal Chemistry* 39, 3929-3937 (1996) whose disclosures are incorporated by reference herein in their entirety.

### **LPS-induced TNF $\alpha$ Production in Mice and Rats**

In order to evaluate *in vivo* inhibition of LPS-induced TNF $\alpha$  production in  
30 rodents, both mice and rats are injected with LPS.

#### Mouse Method

Male Balb/c mice from Charles River Laboratories are pretreated (30 minutes) with compound or vehicle. After the 30 min. pretreat time, the mice are given LPS (lipopolysaccharide from *Escherichia coli* Serotype 055-85, Sigma Chemical Co., St  
35 Louis, MO) 25 ug/mouse in 25 ul phosphate buffered saline (pH 7.0) intraperitoneally. Two hours later the mice are killed by CO<sub>2</sub> inhalation and blood samples are collected by exsanguination into heparinized blood collection tubes and stored on ice. The blood

samples are centrifuged and the plasma collected and stored at  $-20^{\circ}\text{C}$  until assayed for  $\text{TNF}\alpha$  by ELISA.

#### Rat Method

5 Male Lewis rats from Charles River Laboratories are pretreated at various times with compound or vehicle. After a determined pretreat time, the rats are given LPS (lipopolysaccharide from *Escherichia coli* Serotype 055-85, Sigma Chemical Co., St Louis, MO) 3.0 mg/kg intraperitoneally. The rats are killed by  $\text{CO}_2$  inhalation and heparinized whole blood is collected from each rat by cardiac puncture 90 minutes after  
10 the LPS injection. The blood samples are centrifuged and the plasma collected for analysis by ELISA for  $\text{TNF}\alpha$  levels.

#### ELISA Method

$\text{TNF}\alpha$  levels were measured using a sandwich ELISA, as described in Olivera et al., Circ. Shock, 37, 301-306, (1992), whose disclosure is incorporated by reference in  
15 its entirety herein, using a hamster monoclonal antimurine  $\text{TNF}\alpha$  (Genzyme, Boston, MA) as the capture antibody and a polyclonal rabbit antimurine  $\text{TNF}\alpha$  (Genzyme) as the second antibody. For detection, a peroxidase-conjugated goat antirabbit antibody (Pierce, Rockford, IL) was added, followed by a substrate for peroxidase (1 mg/ml orthophenylenediamine with 1% urea peroxide).  $\text{TNF}\alpha$  levels in the plasma samples  
20 from each animal were calculated from a standard curve generated with recombinant murine  $\text{TNF}\alpha$  (Genzyme).

#### **LPS-Stimulated Cytokine Production In Human Whole Blood**

25 **Assay:** Test compound concentrations were prepared at 10 X concentrations and LPS prepared at 1  $\mu\text{g}/\text{mL}$  (final conc. of 50 ng/mL LPS) and added in 50  $\mu\text{L}$  volumes to 1.5 mL eppendorf tubes. Heparinized human whole blood was obtained from healthy volunteers and was dispensed into eppendorf tubes containing compounds and LPS in 0.4 mL volumes and the tubes incubated at 37 C. Following a 4 hour incubation, the  
30 tubes were centrifuged at 5000 rpm for 5 minutes in a TOMY microfuge, plasma was withdrawn and frozen at  $-80^{\circ}\text{C}$ .

Cytokine measurement: IL-1 and/or TNF were quantified using a standardized ELISA technology. An in-house ELISA kit was used to detect human IL-1 and TNF.

35 Concentrations of IL-1 or TNF were determined from standard curves of the appropriate cytokine and  $\text{IC}_{50}$  values for test compound (concentration that inhibited 50% of LPS-stimulated cytokine production) were calculated by linear regression analysis.

**CSBP Kinase Assay:**

This assay measures the CSBP-catalyzed transfer of  $^{32}\text{P}$  from [ $\alpha$ - $^{32}\text{P}$ ]ATP to threonine residue in an epidermal growth factor receptor (EGFR)-derived peptide (T669) with the following sequence: KRELVEPLTPSGEAPNQALLR (residues 661-681). (See Gallagher et al., "Regulation of Stress Induced Cytokine Production by Pyridinyl Imidazoles: Inhibition of CSPB Kinase", BioOrganic & Medicinal Chemistry, to be published 1996).

Kinase reactions (total volume 30  $\mu\text{l}$ ) contain: 25 mM Hepes buffer, pH 7.5; 10 mM  $\text{MgCl}_2$ ; 170  $\mu\text{M}$  ATP<sup>(1)</sup>; 10  $\mu\text{M}$  Na ortho vanadate; 0.4 mM T669 peptide; and 20-80 ng of yeast-expressed purified CSBP2 (see Lee et al., *Nature* 300, n(72), 739-746 (Dec. 1994)). Compounds (5  $\mu\text{l}$  from [6X] stock<sup>(2)</sup>) are pre-incubated with the enzyme and peptide for 20 min. on ice prior to starting the reactions with  $^{32}\text{P}$ /MgATP. Reactions are incubated at 30  $^\circ\text{C}$  for 10 min. and stopped by adding 10  $\mu\text{l}$  of 0.3 M phosphoric acid.  $^{32}\text{P}$ -labeled peptide is separated on phosphocellulose (Wattman, p81) filters by spotting 30  $\mu\text{l}$  reaction mixture. Filters are washed 3 times with 75 mM phosphoric acid followed by 2 washes with  $\text{H}_2\text{O}$ , and counted for  $^{32}\text{P}$ .

(1) The  $K_m$  of CSBP for ATP was determined to be 170  $\mu\text{M}$ . Therefore, compounds screened at the  $K_m$  value of ATP.

(2) Compounds are usually dissolved in DMSO and are diluted in 25 mM HEPES buffer to get final concentration of DMSO of 0.17%.

Representative compounds of Formula (I) and (II), Examples 2, 3, 4, 7 and 8 herein all demonstrated a positive inhibitory activity in this assay having an  $\text{IC}_{50} < 50 \mu\text{M}$ .

**Prostaglandin endoperoxide synthase-2 (PGHS-2) assay:**

This assay describes a method for determining the inhibitory effects of compounds of Formula (I) on human PGHS-2 protein expression in LPS stimulated human monocytes. A suitable assay for PGHS-2 protein expression may be found in a number of publications, including US Patent 5,593,992 whose disclosure is incorporated herein by reference.

**TNF- $\alpha$  in Traumatic Brain Injury Assay**

This assay provides for examination of the expression of tumor necrosis factor mRNA in specific brain regions which follow experimentally induced lateral fluid-percussion traumatic brain injury (TBI) in rats. Since TNF- $\alpha$  is able to induce nerve growth factor (NGF) and stimulate the release of other cytokines from activated

astrocytes, this post-traumatic alteration in gene expression of TNF-  $\alpha$  plays an important role in both the acute and regenerative response to CNS trauma. A suitable assay may be found in WO 97/35856 whose disclosure is incorporated herein by reference.

5

#### **CNS Injury model for IL- $\beta$ mRNA**

This assay characterizes the regional expression of interleukin-1 $\beta$  (IL-1 $\beta$ ) mRNA in specific brain regions following experimental lateral fluid-percussion traumatic brain injury (TBI) in rats. Results from these assays indicate that following  
10 TBI, the temporal expression of IL-1 $\beta$  mRNA is regionally stimulated in specific brain regions. These regional changes in cytokines, such as IL-1 $\beta$  play a role in the post-traumatic pathologic or regenerative sequelae of brain injury. A suitable assay may be found in WO 97/35856 whose disclosure is incorporated herein by reference.

#### **15 Angiogenesis Assay:**

Described below is an assay for determination of inflammatory angiogenesis which may be used to show that cytokine inhibition will stop the tissue destruction of excessive or inappropriate proliferation of blood vessels.

The murine airpouch granuloma model of chronic inflammation (Kimura et al.,  
20 1985, J. Pharmacobio-Dyn., 8:393-400; Colville-Nash et al., 1995, J. Pharm. and Exp. Ther., 274:1463-1472) whose disclosure is incorporated herein by reference in its entirety, is characterized by inflammatory cell influx, fibrous tissue proliferation and intense angiogenesis. It is representative of inflammatory angiogenesis and demonstrates that the angiogenic component can be pharmacologically modulated  
25 independently of granuloma growth and size. In addition, angiogenesis can be accurately quantitated by a vascular casting method.

The effect of a compound on vascular density (and dry weight) is measured for 6 days after induction of the granuloma. This time point has previously been determined to be at or near the peak of angiogenesis. A positive control  
30 medroxyprogesterone, an angiostatic steroid (Gross et al., 1981, Proc. Natl. Acad. Sci. USA, 78:1176-1180) - whose disclosure is hereby incorporated by reference in its entirety, is utilized.

#### **Methods: Murine air pouch granuloma model:**

Day -1, mice are anesthetized using Aerrane (isoflurane) gas (5%), after which  
35 3mls of air is injected into the dorsal subcutaneous tissue using a 27g needle. Mice are allowed to recover.

Day 0, mice are again anesthetized using Aerrane, once anesthetized 0.5ml of Freund's complete adjuvant with 0.1% v/v croton oil is injected into the air pouch formed on Day -1. The animals also begin their dosing regime (number of days dependent upon study) with the animals typically receiving compound in 0.2ml N,N, Dimethyl Acetoacetamide(DMA)(Sigma, St. Louis, Mo.)/Cremephor El (Sigma, St. Louis, Mo.)/ saline (10/10/80) or other appropriate vehicle. The animals are allowed to recover and all subsequent dosing is performed on the animals in the absence of anesthetics.

Days 1-5, animals are dosed according to schedule.

On Day 6 the animals are again anesthetized using Aerrane after which a vascular cast is made (Kimura et al., 1986, J.Pharmacobio-Dyn., 9:442-446), this involves a 1ml tail vein i.v. injection of a Carmine Red(10%)(Sigma, St. Louis, Mo.)/gelatin (5%)(Sigma, St. Louis, Mo.) solution. The animals are then sacrificed by lethal dose of anesthesia and chilled at 4 C for 2 hours prior to the removal of the granuloma tissue.

When the granuloma is removed it is weighed and then dried for 3 days at 45 C and reweighed. The dried tissue is then digested in 0.9ml of a 0.05M phosphate buffer pH 7.0 containing 12 U /ml<sup>-1</sup> papain (Sigma, St. Louis, Mo.) and 0.33g / L<sup>-1</sup> N-acetyl-L-Cysteine (Sigma, St. Louis, Mo.) at 57 C for 3 days. After 3 days digestion the carmine red is solubilized by the addition of 0.1ml 5mM NaOH. Samples are centrifuged and then filtered using 0.2um acrodiscs. The carmine content is then determined against a carmine red standard curve (0.5 to 2 mg/ml) generated in extracted tissue from non carmine treated animals and read at 490nm. Sample and standard values are determined using DeltaSoft Elisa analysis software (Biometallics Inc., Princeton, NJ). The carmine content is then used to determine the vascular indexes for the various treatments, vascular index being the mg carmine dye/gm dry tissue.

Tissue extracts are made by homogenizing granulomas in 0.5ml 5mM KH<sub>2</sub>PO<sub>4</sub> / 0.1gm wet tissue. IL-1 $\beta$  levels are determined using a Cytoscreen Immunoassay Kit (catalog # KMC 0012) from BioSource International, Camarillo, CA. TNF- $\alpha$  levels are determined using the following assay: plates were coated with hamster anti-murine TNF- $\alpha$  antibody (Genzyme, Cambridge, MA), for 2 hours at 37°C, washed and blocked with a casein-BSA solution (5gram/L for each) for 1 hour at 37°C, the samples are added and incubated at 4°C overnight. Plates are washed, and the secondary antibody, Rabbit Anti-mouse TNF- $\alpha$  (Genzyme), was added for 2 hours at 37°C, the plates were washed, and the tertiary antibody Goat Anti-rabbit peroxidase conjugate (BioSource International, Camarillo, CA) is added for 2 hours at 37°C. The plates are then washed, and substrate OPD (Sigma) is added for 20 minutes at room temperature. The reaction




is terminated with 25ul 0.1M NaF per well, the O.D. read at 460nm. Sample values for both ELISAs are calculated using DeltaSoft ELISA analysis software (Biomettalics Inc., Princeton, NJ).

5           All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

10           The above description fully discloses the invention including preferred embodiments thereof. Modifications and improvements of the embodiments specifically disclosed herein are within the scope of the following claims. Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. Therefore the Examples herein are to be construed as merely illustrative and not a limitation of the scope of the  
15           present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.

What is Claimed Is:

1. A compound of the formula



(I)

- 5 wherein

R<sub>1</sub> is 4-pyridyl, 4-pyrimidinyl, 4-pyridazinyl, 1,2,4-triazin-5-yl, 4-quinolyl, 6-isoquinolyl, quinazolin-4-yl, 1-imidazolyl or 1-benzimidazolyl ring, which ring is optionally substituted independently one to three times with Y, NHR<sub>a</sub>, optionally substituted C<sub>1-4</sub> alkyl, halogen, hydroxyl, optionally substituted C<sub>1-4</sub> alkoxy, optionally substituted C<sub>1-4</sub> alkylthio, optionally substituted C<sub>1-4</sub> alkylsulfinyl, CH<sub>2</sub>OR<sub>12</sub>, amino, mono and di- C<sub>1-6</sub> alkyl substituted amino, N(R<sub>10</sub>)C(O)R<sub>b</sub>, N(R<sub>10</sub>)S(O)<sub>2</sub>R<sub>d</sub>, or an N-heterocyclyl ring which ring has from 5 to 7 members and optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>15</sub>;

Y is X<sub>1</sub>-R<sub>a</sub>;

- 15 X<sub>1</sub> is sulfur or oxygen;

R<sub>a</sub> is C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic, heterocyclylC<sub>1-6</sub> alkyl, heteroaryl, or heteroarylC<sub>1-6</sub>alkyl, wherein each of these moieties may be optionally substituted;

- 20 R<sub>b</sub> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl;

R<sub>d</sub> is C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl;

- 25 R<sub>4</sub> is phenyl, naphth-1-yl or naphth-2-yl, or heteroaryl, which is optionally substituted by one to three substituents, each of which is independently selected, and which, for a 4-phenyl, 4-naphth-1-yl, 5-naphth-2-yl or 6-naphth-2-yl substituent, is halogen, cyano, nitro, C(Z)NR<sub>7</sub>R<sub>17</sub>, C(Z)OR<sub>16</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>COR<sub>12</sub>, SR<sub>5</sub>, S(O)R<sub>5</sub>, OR<sub>12</sub>, halo-substituted-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl, ZC(Z)R<sub>12</sub>, NR<sub>10</sub>C(Z)R<sub>16</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>10</sub>R<sub>20</sub> and which, for other positions of substitution, is halogen, cyano, nitro, phenyl, C(Z)NR<sub>13</sub>R<sub>14</sub>, C(Z)OR<sub>3</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>m</sub>"COR<sub>3</sub>, S(O)<sub>m</sub>R<sub>3</sub>, OR<sub>3</sub>, halo-substituted-C<sub>1-4</sub> alkyl, C<sub>1-10</sub> alkyl, ZC(Z)R<sub>3</sub>, optionally substituted phenyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>m</sub>"NR<sub>10</sub>C(Z)R<sub>3</sub>, NR<sub>10</sub>S(O)<sub>m</sub>'R<sub>8</sub>, NR<sub>10</sub>S(O)<sub>m</sub>'NR<sub>7</sub>R<sub>17</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>m</sub>"NR<sub>13</sub>R<sub>14</sub>;

m is 0, or the integer 1 or 2;

m' is an integer having a value of 1 or 2,


- m" is 0, or an integer having a value of 1 to 5;  
 n is 0, or an integer having a value of 1 to 10;  
 v is 0, or an integer having a value of 1 or 2;  
 R<sub>2</sub> is hydrogen, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub> OR<sub>9</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OR<sub>11</sub>, C<sub>1-10</sub>alkyl, halo-substituted  
 5 C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub>  
 alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl,  
 heteroaryl, heteroarylC<sub>1-10</sub>alkyl, heterocyclyl, heterocyclylC<sub>1-10</sub> alkyl,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>13</sub>R<sub>14</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NO<sub>2</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>S(O)<sub>m</sub>'NR<sub>13</sub>R<sub>14</sub>,  
 10 (CR<sub>10</sub>R<sub>230</sub>)<sub>n</sub>C(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OC(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)OR<sub>11</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)NR<sub>11</sub>OR<sub>9</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>N(OR<sub>6</sub>)C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>N(OR<sub>6</sub>)C(Z)R<sub>11</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(=NOR<sub>6</sub>)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>19</sub>)NR<sub>13</sub>R<sub>14</sub>,  
 15 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OC(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>,  
 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>10</sub>, 5-(R<sub>18</sub>)-1,2,4-oxadiazol-3-yl or 4-(R<sub>12</sub>)-5-(R<sub>18</sub>R<sub>19</sub>)-  
 4,5-dihydro-1,2,4-oxadiazol-3-yl; wherein the cycloalkyl, cycloalkyl alkyl, aryl,  
 arylalkyl, heteroaryl, heteroaryl alkyl, heterocyclic and heterocyclic alkyl groups  
 may be optionally substituted;  
 20 R<sub>3</sub> is heterocyclyl, heterocyclylC<sub>1-10</sub> alkyl or R<sub>8</sub>;  
 R<sub>5</sub> is hydrogen, C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl or NR<sub>7</sub>R<sub>17</sub>, excluding the  
 moieties SR<sub>5</sub> being SNR<sub>7</sub>R<sub>17</sub> and SOR<sub>5</sub> being -SOH;  
 R<sub>6</sub> is hydrogen, a pharmaceutically acceptable cation, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl,  
 aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclic, aroyl, or C<sub>1-10</sub>  
 25 alkanoyl;  
 R<sub>7</sub> and R<sub>17</sub> is each independently selected from hydrogen or C<sub>1-4</sub> alkyl or R<sub>7</sub> and R<sub>17</sub>  
 together with the nitrogen to which they are attached form a heterocyclic ring of 5  
 to 7 members which ring optionally contains an additional heteroatom selected from  
 oxygen, sulfur or NR<sub>15</sub>;  
 30 R<sub>8</sub> is C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub>  
 cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub>  
 alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>11</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>18</sub>,  
 (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>13</sub>R<sub>14</sub>; wherein the aryl, arylalkyl, heteroaryl, heteroaryl alkyl may  
 be optionally substituted;  
 35 R<sub>9</sub> is hydrogen, C(Z)R<sub>11</sub> or optionally substituted C<sub>1-10</sub> alkyl, S(O)<sub>2</sub>R<sub>18</sub>, optionally  
 substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl;  
 R<sub>10</sub> and R<sub>20</sub> is each independently selected from hydrogen or C<sub>1-4</sub> alkyl;

- R<sub>11</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or heteroarylC<sub>1-10</sub> alkyl, wherein the aryl, arylalkyl, heteroaryl, heteroaryl alkyl, heterocyclyl or heterocyclylalkyl, may be optionally substituted;
- 5 R<sub>12</sub> is hydrogen or R<sub>16</sub>;
- R<sub>13</sub> and R<sub>14</sub> is each independently selected from hydrogen or optionally substituted C<sub>1-4</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl, or together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from
- 10 oxygen, sulfur or NR<sub>9</sub> ;
- R<sub>15</sub> is hydrogen, C<sub>1-4</sub> alkyl or C(Z)-C<sub>1-4</sub> alkyl;
- R<sub>16</sub> is C<sub>1-4</sub> alkyl, halo-substituted-C<sub>1-4</sub> alkyl, or C<sub>3-7</sub> cycloalkyl;
- R<sub>18</sub> is C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, aryl, arylC<sub>1-10</sub> alkyl, heterocyclyl, heterocyclyl-C<sub>1-10</sub>alkyl, heteroaryl or heteroarylalkyl, wherein the aryl, arylalkyl,
- 15 heteroaryl, heteroaryl alkyl, heterocyclyl or heterocyclylalkyl may be optionally substituted;
- R<sub>19</sub> is hydrogen, cyano, C<sub>1-4</sub> alkyl, C<sub>3-7</sub> cycloalkyl or aryl;
- R<sub>23</sub> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl moiety, all of which
- 20 may be optionally substituted;
- Z is oxygen or sulfur;
- provided that when R<sub>1</sub> is a 4-pyridyl, R<sub>4</sub> is a 4-NH<sub>2</sub>S(O)<sub>2</sub>phenyl, then R<sub>2</sub> is other than a 3-position trifluoromethyl group;
- or a pharmaceutically acceptable salt thereof.
- 25
2. The compound according to Claim 1 wherein R<sub>1</sub> is a substituted 4-pyrimidinyl.
3. The compound according to Claim 2 wherein the substituent is Y or NHR<sub>a</sub>.
- 30 4. The compound according to Claim 1 wherein R<sub>1</sub> is a substituted 4-pyridyl.
5. The compound according to Claim 4 wherein the substituent is Y or NHR<sub>a</sub>.
6. The compound according to Claim 2 or 4 wherein R<sub>4</sub> is an optionally
- 35 substituted phenyl.

7. The compound according to Claim 4 wherein the phenyl is substituted one or more times independently by halogen, SR<sub>5</sub>, S(O)R<sub>5</sub>, OR<sub>12</sub>, halo-substituted-C<sub>1-4</sub> alkyl, or C<sub>1-4</sub> alkyl.
- 5 8. The compound according to Claim 1 wherein R<sub>2</sub> is selected from optionally substituted C<sub>4</sub> to C<sub>6</sub> cycloalkyl, optionally substituted C<sub>4</sub> or C<sub>6</sub> cycloalkyl C<sub>1-4</sub> alkyl, optionally substituted heterocyclic, optionally substituted heterocyclic C<sub>1-4</sub> alkyl, optionally substituted aryl, or an optionally substituted aryl C<sub>1-4</sub> alkyl.
- 10 9. The compound according to Claim 1 which is:  
4-[1-(4-Fluorophenyl)-3-phenyl-1*H*-pyrazol-5-yl]pyridine  
4-[4-Bromo-1-(4-fluorophenyl)-3-phenyl-1*H*-pyrazol-5-yl]pyridine  
4-[1-(4-Fluorophenyl)-3-[4-(methylthio)phenyl]-1*H*-pyrazol-5-yl]pyridine  
4-[1-(4-Fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-1*H*-pyrazol-5-yl]pyridine  
15 4-[1-(4-Fluorophenyl)-3-[4-(methylsulfinyl)phenyl]-1*H*-pyrazol-5-yl]pyridine; or a pharmaceutically acceptable salt thereof.
10. A pharmaceutical composition comprising a compound according to any of Claims 1 to 9 and a pharmaceutically acceptable carrier or diluent.
- 20 11. A method of treating a CSBP/RK/p38 kinase mediated disease, in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of Formula (I) according to any of Claims 1 to 9.
- 25 12. The method according to Claim 11 wherein the mammal is afflicted with a CSBP/RK/p38 kinase mediated disease which is psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis and acute synovitis, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other  
30 arthritic condition, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, Alzheimer's disease, stroke, neurotrauma, asthma, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcososis, bone resorption disease, osteoporosis, restenosis, cardiac and renal reperfusion injury, chronic renal failure, congestive heart failure, angiogenic  
35 diseases, cancer, thrombosis, glomerulonephritis, diabetes, graft vs. host reaction, allograft rejection, inflammatory bowel disease, Crohn's disease, ulcerative colitis, multiple sclerosis, muscle degeneration, diabetic retinopathy, macular degeneration,

tumor growth and metastasis, eczema, contact dermatitis, psoriasis, sunburn, and conjunctivitis.

13. A compound of the formula



(II)

wherein

R<sub>1</sub> is 4-pyridyl, 4-pyrimidinyl, 4-pyridazinyl, 4-triazin-5-yl, quinolyl, isoquinolyl, quinazolin-4-yl, 1-imidazolyl or 1-benzimidazolyl ring, which ring is optionally substituted independently one to three times with Y, NHR<sub>a</sub>, optionally substituted C<sub>1-4</sub> alkyl, halogen, hydroxyl, optionally substituted C<sub>1-4</sub> alkoxy, optionally substituted C<sub>1-4</sub> alkylthio, optionally substituted C<sub>1-4</sub> alkylsulfinyl, CH<sub>2</sub>OR<sub>12</sub>, amino, mono and di- C<sub>1-6</sub> alkyl substituted amino, N(R<sub>10</sub>)C(O)R<sub>b</sub>; N(R<sub>10</sub>)S(O)<sub>2</sub>R<sub>d</sub> or an N-heterocyclyl ring which ring has from 5 to 7 members and optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>15</sub>;

Y is X<sub>1</sub>-R<sub>a</sub>;

X<sub>1</sub> is sulfur or oxygen;

R<sub>a</sub> is C<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, heterocyclic, heterocyclylC<sub>1-6</sub> alkyl, heteroaryl, or heteroarylC<sub>1-6</sub>alkyl, wherein each of these moieties may be optionally substituted;

R<sub>b</sub> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl;

R<sub>d</sub> is C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl;

R<sub>4</sub> is phenyl, naphth-1-yl or naphth-2-yl, a fused phenyl containing ring system, or a heteroaryl, which is optionally substituted by one to three substituents, each of which is independently selected, and which, for a 4-phenyl, 4-naphth-1-yl, 5-naphth-2-yl or 6-naphth-2-yl substituent, is halogen, cyano, nitro, C(Z)NR<sub>7</sub>R<sub>17</sub>, C(Z)OR<sub>16</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>COR<sub>12</sub>, SR<sub>5</sub>, SOR<sub>5</sub>, OR<sub>12</sub>, halo-substituted-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl, ZC(Z)R<sub>12</sub>, NR<sub>10</sub>C(Z)R<sub>16</sub>, or (CR<sub>10</sub>R<sub>20</sub>)<sub>v</sub>NR<sub>10</sub>R<sub>20</sub> and which, for other positions of substitution, is halogen, cyano, C(Z)NR<sub>13</sub>R<sub>14</sub>, C(Z)OR<sub>3</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>m</sub>COR<sub>3</sub>, S(O)<sub>m</sub>R<sub>3</sub>, OR<sub>3</sub>, halo-substituted-C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkyl,

- (CR<sub>10</sub>R<sub>20</sub>)<sub>m</sub>"NR<sub>10</sub>C(Z)R<sub>3</sub>, NR<sub>10</sub>S(O)<sub>m</sub>R<sub>8</sub>, NR<sub>10</sub>S(O)<sub>m</sub>'NR<sub>7</sub>R<sub>17</sub>, ZC(Z)R<sub>3</sub> or (CR<sub>10</sub>R<sub>20</sub>)<sub>m</sub>"NR<sub>13</sub>R<sub>14</sub>;
- n is 0, or an integer having a value of 1 to 10;
- v is 0, or an integer having a value of 1 or 2;
- 5 m is 0, or the integer 1 or 2;
- m' is an integer having a value of 1 or 2,
- m" is 0, or an integer having a value of 1 to 5;
- Z is oxygen or sulfur;
- R<sub>2</sub> is hydrogen, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub> OR<sub>9</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OR<sub>11</sub>, C<sub>1-10</sub>alkyl, halo-substituted
- 10 C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-10</sub> alkyl, C<sub>5-7</sub> cycloalkenyl, C<sub>5-7</sub> cycloalkenyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub>alkyl, heterocyclyl, heterocyclylC<sub>1-10</sub> alkyl, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NO<sub>2</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>CN, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>S(O)<sub>m</sub>'NR<sub>13</sub>R<sub>14</sub>,
- 15 (CR<sub>10</sub>R<sub>230</sub>)<sub>n</sub>C(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OC(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)OR<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(Z)NR<sub>11</sub>OR<sub>9</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>N(OR<sub>6</sub>)C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>N(OR<sub>6</sub>)C(Z)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>C(=NOR<sub>6</sub>)R<sub>11</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(=NR<sub>19</sub>)NR<sub>13</sub>R<sub>14</sub>,
- 20 (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>OC(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)NR<sub>13</sub>R<sub>14</sub>, (CR<sub>10</sub>R<sub>23</sub>)<sub>n</sub>NR<sub>10</sub>C(Z)OR<sub>10</sub>, 5-(R<sub>18</sub>)-1,2,4-oxadiazol-3-yl or 4-(R<sub>12</sub>)-5-(R<sub>18</sub>R<sub>19</sub>)-4,5-dihydro-1,2,4-oxadiazol-3-yl; wherein the cycloalkyl, cycloalkyl alkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclic and heterocyclic alkyl groups may be optionally substituted;
- 25 R<sub>3</sub> is heterocyclyl, heterocyclylC<sub>1-10</sub> alkyl or R<sub>8</sub>;
- R<sub>5</sub> is hydrogen, C<sub>1-4</sub> alkyl, C<sub>2-4</sub> alkenyl, C<sub>2-4</sub> alkynyl or NR<sub>7</sub>R<sub>17</sub>, excluding the moieties SR<sub>5</sub> being SNR<sub>7</sub>R<sub>17</sub> and SOR<sub>5</sub> being SOH;
- R<sub>6</sub> is hydrogen, a pharmaceutically acceptable cation, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl, heteroarylC<sub>1-4</sub> alkyl, heterocyclic, aroyl, or C<sub>1-10</sub>
- 30 alkanoyl;
- R<sub>7</sub> and R<sub>17</sub> is each independently selected from hydrogen or C<sub>1-4</sub> alkyl or R<sub>7</sub> and R<sub>17</sub> together with the nitrogen to which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>15</sub>;
- 35 R<sub>8</sub> is C<sub>1-10</sub> alkyl, halo-substituted C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl, heteroarylC<sub>1-10</sub> alkyl, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>OR<sub>11</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>S(O)<sub>m</sub>R<sub>18</sub>, (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NHS(O)<sub>2</sub>R<sub>18</sub>,

- (CR<sub>10</sub>R<sub>20</sub>)<sub>n</sub>NR<sub>13</sub>R<sub>14</sub>; wherein the aryl, arylalkyl, heteroaryl, heteroaryl alkyl may be optionally substituted;
- R<sub>9</sub> is hydrogen, -C(Z)R<sub>11</sub> or optionally substituted C<sub>1-10</sub> alkyl, S(O)<sub>2</sub>R<sub>18</sub>, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl;
- 5 R<sub>10</sub> and R<sub>20</sub> is each independently selected from hydrogen or C<sub>1-4</sub> alkyl;
- R<sub>11</sub> is hydrogen, C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, heterocyclyl C<sub>1-10</sub>alkyl, aryl, arylC<sub>1-10</sub> alkyl, heteroaryl or heteroarylC<sub>1-10</sub> alkyl;
- R<sub>12</sub> is hydrogen or R<sub>16</sub>;
- R<sub>13</sub> and R<sub>14</sub> is each independently selected from hydrogen or optionally substituted
- 10 C<sub>1-4</sub> alkyl, optionally substituted aryl or optionally substituted aryl-C<sub>1-4</sub> alkyl, or together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR<sub>9</sub> ;
- R<sub>15</sub> is hydrogen, C<sub>1-4</sub> alkyl or C(Z)-C<sub>1-4</sub> alkyl;
- 15 R<sub>16</sub> is C<sub>1-4</sub> alkyl, halo-substituted-C<sub>1-4</sub> alkyl, or C<sub>3-7</sub> cycloalkyl;
- R<sub>18</sub> is C<sub>1-10</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclyl, aryl, arylC<sub>1-10</sub> alkyl, heterocyclyl, heterocyclyl-C<sub>1-10</sub>alkyl, heteroaryl or heteroarylalkyl;
- R<sub>19</sub> is hydrogen, cyano, C<sub>1-4</sub> alkyl, C<sub>3-7</sub> cycloalkyl or aryl;
- R<sub>23</sub> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> cycloalkyl, aryl, arylC<sub>1-4</sub> alkyl, heteroaryl,
- 20 heteroarylC<sub>1-4</sub>alkyl, heterocyclyl, or heterocyclylC<sub>1-4</sub> alkyl moiety, all of which may be optionally substituted;
- or a pharmaceutically acceptable salt thereof.
14. The compound according to Claim 13 wherein R<sub>1</sub> is a substituted 4-pyridyl or
- 25 4-pyrimidinyl.
15. The compound according to Claim 14 wherein the substituent is Y, or NHR<sub>a</sub>.
16. The compound according to Claim 14 wherein R<sub>4</sub> is an optionally substituted
- 30 phenyl.
17. The compound according to Claim 16 wherein the phenyl is substituted one or more times independently by halogen, -SR<sub>5</sub>, S(O)R<sub>5</sub>, OR<sub>12</sub>, halo-substituted-C<sub>1-4</sub> alkyl, or C<sub>1-4</sub> alkyl.
- 35 18. The compound according to Claim 13 wherein R<sub>2</sub> is selected from optionally substituted C<sub>4</sub> to C<sub>6</sub>cycloalkyl, optionally substituted C<sub>4</sub> or C<sub>6</sub> cycloalkylC<sub>1-4</sub> alkyl,



optionally substituted heterocyclic, optionally substituted heterocyclicalkyl, optionally substituted aryl, or optionally substituted aryl alkyl.

19. The compound according to Claim 13 which is:


- 5 4-[1-(4-Fluorophenyl)-4,5-dihydro-3-phenyl-1*H*-pyrazol-5-yl]pyridine  
4-[1-(4-Fluorophenyl)-4,5-dihydro-3-[4-(methylthio)phenyl]-1*H*-pyrazol-5-yl]pyridine,  
or a pharmaceutically acceptable salt thereof.

20. A pharmaceutical composition comprising a compound according to any of  
10 Claims 13 to 19 and a pharmaceutically acceptable carrier or diluent.

21. A method of treating a CSBP/RK/p38 kinase mediated disease, in a mammal in  
need thereof, which comprises administering to said mammal an effective amount of a  
compound of Formula (I) according to any of Claims 13 to 19.

- 15 22. The method according to claim 21 wherein the mammal is afflicted with a  
CSBP/RK/p38 kinase mediated disease which is psoriatic arthritis, Reiter's syndrome,  
rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis and acute synovitis,  
rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other  
20 arthritic condition, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic  
shock syndrome, Alzheimer's disease, stroke, neurotrauma, asthma, adult respiratory  
distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis,  
pulmonary sarcososis, bone resorption disease, osteoporosis, restenosis, cardiac and  
renal reperfusion injury, chronic renal failure, congestive heart failure, angiogenic  
25 diseases, cancer, thrombosis, glomerulonephritis, diabetes, graft vs. host reaction,  
allograft rejection, inflammatory bowel disease, Crohn's disease, ulcerative colitis,  
multiple sclerosis, muscle degeneration, diabetic retinopathy, macular degeneration,  
tumor growth and metastasis, eczema, contact dermatitis, psoriasis, sunburn, and  
conjunctivitis.


- 30 23. A process for producing a compound of Formula (I), according to Claim 1 wherein  
R1 is an optionally substituted pyrimidinyl, which process comprises cyclizing a  
compound of the formula



wherein  $R_2$  is an optionally substituted phenyl, as defined according to formula (I);  
with a compound of the formula:  $R_4NHNH_2$ , wherein  $R_4$  is as defined for Formula (I),  
to yield a compound of Formula (I), or if necessary, converting a precursor of  $R_1$ ,  $R_2$   
and  $R_4$  to a group  $R_1$ ,  $R_2$  and  $R_4$ .

5

24. A process for producing a compound of Formula (I), according to Claim 1,  
wherein  $R_1$  is an optionally substituted pyridyl, which process comprises cyclizing a  
compound of the formula



10 wherein  $R_2$  is an optionally substituted phenyl, as defined according to formula (I);  
with a compound of the formula:  $R_4NHNH_2$ , wherein  $R_4$  is as defined for Formula (I),  
to yield a compound of Formula (I), or if necessary, converting a precursor of  $R_1$ ,  $R_2$   
and  $R_4$  to a group  $R_1$ ,  $R_2$  and  $R_4$ .

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/12387

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :A61K 31/44; C07D 401/14, 405/14, 409/14

US CL :514/341; 548/275.4

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/341; 548/275.4

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
Please See Extra Sheet.

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3,678,063 A (BINON et al.) 18 July 1972, entire document.	1-10, 13-20
A	US 5,516,907 A (TALLEY et al.) 14 May 1996, entire document.	1-10, 13-20
A	US 5,580,985 A (LEE et al.) 03 December 1996, entire document.	1-10, 13-20



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

11 AUGUST 1998

Date of mailing of the international search report

08 OCT 1998

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

FLOYD D. HIGEL aco

Telephone No. (703) 308-1235

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/12387

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☒ Claims Nos.: 11, 12  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/12387

## B. FIELDS SEARCHED

Documentation other than minimum documentation that are included in the fields searched:

Chemical Abstracts

Index Chemicus

Current Abstracts of Chemistry